# Calcito

## High efficiency rhizosphere corrector

CALCITO is a rhizosphere corrector based on short-chain carboxylic acids containing calcium and magnesium. CALCIUM thanks to the long-acting organic acidification of the rhizosphere improves the absorption of Nutrients, in particular calcium, and improves the fertility (structure and pH) of saline, sodic and calcareous soils. CALCITO prevents and treats calcium deficiencies (apical rot, cracking, bitter pit, tip burn) and strengthens plant tissues.

**Benefits**

- More balanced and resistant plants (biotic and abiotic stress) thanks to an optimal calcium absorption;
- Better absorption of Nutrients, thank to the long-acting organic acidification of the rhizosphere;
- More capillary rooting and buffering of soil salinity;
- Prevents calcium deficiency diseases: bitter pit, blossom end rot, lettuce hemming;
- More fertile and productive soils as it favors the development of useful microorganisms.

**TYPE**
Liquid

**PACKAGING**





**COMPOSITION**

| | |
|---|---|
| **Calcium ($CaO$) soluble in water** | **9%** |
| **Magnesium (MgO) soluble in water** | **1%** |
| **pH** | **1** |

**DENSITY**
1,25 +/- 0,05 kg/dm$^3$

It also contains low molecular weight carboxylic acids.
Do not mix with products containing phosphorus. Color and density are indicative. For hazard warnings see page 154..

## DOSAGES AND USES

| CROPS | DOSAGES FERTIGATION | PERIOD |
|---|---|---|
| Stone fruits | 10-15 l/ha | preflowering, post fruit set, stone swelling |
| Pome fruits and kiwi | 10-20 l/ha | flowering, after fruit set, fruit swelling |
| Wine and table grape | 10-15 l/ha | vegetative development, fruit set, berry development |
| Citrus trees | 10-15 l/ha | flowering, after fruit set,fruit swelling |
| Open field fruit vegetables | 10-15 l/ha | after 10-15 days from the transplant, repeat every 20 days |
| Open field Leafy vegetables | 10 l/ha | after 7-10 days from the transplant, repeat every 7-10 days |
| Horticultural in greenhouse | 1-2 l/1000 m$^2$ | after 7-10 days from the transplant, repeat every 7-10 days |