



<sup>®</sup>  
C E R E A  
F C P

# PRODUCTS CATALOGUE



Grow well  
to **LIVE** better



# INDEX

## INTRODUCTION

THE COMPANY .....	4
HISTORICAL NOTES .....	6
LEGEND .....	11
CERTIFICATIONS .....	13

## SPECIAL NUTRITION 14

NUTRITIONAL SOLUTIONS .....	16
GUIDELINES .....	18

### NANO.T® - TECHNOLOGY FOR AGRICULTURE 24

NANO.T®Fe .....	30
NANO.T®Fe Bio .....	31
NANO.T®Cu .....	32
NANO.T®Cu Bio .....	33
NANO.T®Total .....	34
NANO.T®Carbo .....	36
NANO.T®CaPO .....	37
NANO.T®Zn .....	38
NANO.T®Chitosano .....	39

### REACTIVE - THE MESO & MICROELEMENTS FORCE 40

Proser MnZn .....	41
Proser Ca .....	42
Calcito .....	43
CalcioMagno .....	44
Color MgZn .....	45
Febo Mix .....	46
Febo Bio .....	47
MagnetiCal .....	48
Focus Ca .....	49
Oro-B .....	50

### FUTURA - BIOSTIMULATING ACTION BY NATURE 52

Verv Plus .....	53
Verv N9 .....	54
Glycos Plus .....	55
StimUp .....	56
Cerere .....	57
B-Power .....	58
K-Fast .....	59
Edafos .....	60
Be-Start 5.15 .....	61
Iride .....	62
Giove Bio N8 .....	63
NaturBlack .....	64
Giove Bio Gold .....	65
SuprEmo .....	66
Crisco .....	67
VigorGreen .....	68
Mycovix .....	70
A-Myco .....	71
Edafix .....	72

### LEAF - MACRO & MESOELEMENTS 74

Leaf N .....	75
Leaf P-Ca .....	76
Leaf K .....	77
Leaf N-Fast .....	78
Leaf S-Quality .....	79
HydroStar BTC .....	80

## GRANULARS 82

THE IMPORTANCE OF PHOSPHORUS .....	84
THE EFFICIENCY OF FERTILIZERS .....	85

BIOACTIVATED FERTILIZERS .....	85
--------------------------------	----

## NUECR4 - THE EFFICIENT FERTILIZER 86

Blurain .....	88
Control .....	89

## POWER - MICROGRANULARS: STARTER EFFECT 90

SuperPower .....	91
SuperPower Humic .....	92
SuperPower Plus .....	93
SuperPower Extra .....	94
Power BioAger .....	95
Power BioMaster .....	96
Power BioNascor .....	97

## FERT - THE BENEFITS OF THE TWO WORLDS 98

Granoro .....	100
Flex .....	101
Ortofrutto Special .....	102
Ortofrutto Special Top .....	103
Master Plus .....	105
Olivo .....	106
Dual Band .....	107
TrioStart .....	108

## FERT PREMIUM - THE BENEFITS OF THE TWO WORLDS 109

Vinfrutto .....	109
Ortoplus .....	110
Jolly .....	111
Vinfrutto Star .....	112

## ACTIVE - LOVE FOR YOUR CROP 114

Cereaphos .....	115
Universal Up .....	116
Land Plus .....	118
Cerea Blu .....	119
Super Red .....	120
Terra 33 .....	121

## ACTIVE PREMIUM - LOVE FOR YOUR CROP 122

Bluactive .....	122
Global .....	123
Red Ball .....	124
Base .....	125
Mastercote .....	126

## ORGANIC - ECO FRIENDLY AGRICULTURE 128

BioNascor .....	129
BioMaster .....	130
BioAger .....	131
Biorganic N11 .....	132
Biocere 533 .....	133

## SPECIALITY - THE FUTURE IS HERE 134

Evolution 56 .....	136
CereaS 38 .....	137
CereaSlow 33 .....	139
CereaSlow 40 .....	140
CereaSlow 46 .....	141
StarSlow .....	142
Più Sprint .....	143

## WATER-SOLUBLE 144

### FERTIGATION - WATER-SOLUBLE FERTILIZER 146

Fertigation 20.20.20 .....	147
Fertigation 10.40.10 .....	148
Fertigation 15.5.26 .....	149
Fertigation 17.6.21 .....	150
Fertigation 7.15.30 .....	151
Fertigation 30.10.10 .....	152

HAZARD WARNINGS .....	154
-----------------------	-----

# THE COMPANY



# FOR OVER 100 YEARS TOGETHER WITH FARMERS.

Our mission  
is clear:  
to feed the plants,  
respecting the  
environment and  
guaranteeing  
quantity, quality  
and health  
of agricultural  
production.

## Where we are, how we work

Our headquarter is located in Bonavicina di San Pietro di Morubio (province of Verona), in the heart of the Po Valley, one of the most fertile and strategic agricultural areas in Italy. Thanks to our logistical location close to major roads, we can guarantee fast transportation, low transport costs, and reliable distribution.

All key production phases are carried out at our facility:

- research and experimental development;
- production of complex granular fertilizers (NPK, NP, PK), mineral and organo-mineral fertilizers;
- customized blending with double screening of granular products;
- production of special formulations for fertigation and foliar applications;
- granule coating;
- quality control.

All systems are computerized and automated to ensure maximum flexibility and promptness in meeting customer requests.

## Vision

Together with the farmer for sustainable agriculture.

## Our values

Integrity, Transparency, Quality, Innovation, Passion, Mutuality.

# HISTORICAL NOTES

## MORE THAN A CENTURY OF INNOVATION.


Our history isn't just a list of milestones, but a demonstration of our ongoing commitment: innovating without forgetting our origins.

Today, as in 1908, we remain close to farmers, capable of listening, to offer effective solutions and look with confidence toward the most important challenge: sustainable agriculture for future generations.



**2014**

**QUALITY CONTROL, RESEARCH, EXPERIMENTATION**  
Establishment of an agronomic laboratory with a growth chamber for R&D activities. Renovation of the chemical laboratory for quality control of fertilizers and raw materials




**2008**

**SOLAR SYSTEM**  
Construction of a photovoltaic system on the factory roof (0.8 MW capacity)




**2002**

**ORGANO-MINERAL FERTILIZERS**  
Beginning of organo-mineral fertilizer production



**2018**

**COATING PLANT**  
Construction of a plant to enrich each granular product with both liquid and powdered noble materials.



**2020**

**NEW TECHNOLOGY**  
CEREA FCP obtains the patent for the NANO.T technology which allows to produce highly efficient nanofertilizers.



**2021**

**SUSTAINABILITY AND SAFETY**  
The company is at the forefront of the circular economy, as it is committed to using raw materials from renewable sources.



**1908**

**THE FACTORY**  
Fabbrica Cooperativa  
Perfosfati was founded



**1910**

**SULFURIC ACID AND  
SUPERPHOSPHATE**  
First production of sulfuric  
acid and superphosphate



**1990**

**NEW FACTORY**  
New office and  
factory: from Cerea to  
Bonavicina.




**1977**

**MIXING SYSTEM**  
New machine for the production  
of mixed compounds fertilizers,  
among the first in Italy..  
CEREA FCP can now produce  
tailor-made products according to  
customers' needs.



**1952**

**NEW FERTILIZERS**  
Start of granular  
fertilizer production.



**2023**

**NEW PLANT**  
Expansion and upgrading,  
according to the canons of  
Industry 4.0, of the plant for the  
production of liquid fertilizers  
to be applied in fertigation and  
foliar application



**2025**

**NEW LABORATORIES**  
Construction of new laboratories  
dedicated to Research and Development  
activities, including powder grinding  
rooms, instrumentation rooms, R&D,  
agronomy, and a microbiology laboratory.

# CEREA FCP TODAY

Cerea FCP is a **cooperative founded by farmers** in 1908 that, just as it did back then, still produces fertilizers today for its members and for farmers around the world.

The company has **laboratories for quality control** of raw materials and fertilizers. Regular **inspections and audits are conducted to ensure product quality** before it reaches the customer.



**New products are tested through experimental trials** and subsequently by our collaborators and trusted customers.

During 2025, CEREA FCP **renovated its laboratories** with innovative equipment. These facilities are dedicated to analysis, R&D, and agronomy.

Cerea FCP is able **to create specific formulations** to meet the needs of individual farmers.

CEREA FCP has always been committed to technological innovation applied to plant nutrition. This ongoing dedication has enabled the **development and patenting of NANO.T technology** which today sets a new standard in terms of nutritional efficiency and sustainability.

CEREA FCP has also **patented a technique for plant biofortification** (increasing certain nutrients to ensure a **healthier and more complete diet for consumers**).



Today, the company's products (granular, water-soluble, and liquid for specialty Nutrition) are available **throughout Italy**. The company exports **to Europe and many other countries around the world**.

The decision to produce **renewable energy** (installation of an 800 kW photovoltaic system), the concrete application of the circular economy and the adoption of an energy efficiency plan have led Cerea FCP to improve the company's ecological footprint.



**www.fcpcerea.it**  
**www.nanot.eu**

The company's websites aim to serve as a resource for anyone seeking information about cultivation techniques, products, research projects, and other Cerea FCP initiatives.

The nanot.eu website is aimed specifically at anyone interested in learning more about nanoNutrients.

The website is a tool for spreading awareness of the company's values and philosophy, and at the same time, an effective way to inform and communicate with all possible audiences, whether retailers, farmers, agronomists, researchers, or simply enthusiasts.



**FOLLOW US**

---

## **Contacts**

### **Customer service/ Italy**

Phone +39 045 7125914

### **Customer service/ Abroad**

Phone +39 045 7125916

### **Agronomic service**

agronomia@fcpcerea.it

### **General company number**

Phone +39 045 7125911

### **E-mail**

fcpcerea@fcpcerea.it

### **Headquarter**

**FABBRICA COOPERATIVA PERFOSFATI CERIA**

Via Farfusola 6, 37050

Bonavicina di S.Pietro di Morubio (VR)





# LEGEND

## SYMBOLS

---



Phosphorus, expressed in label as  $P_2O_5$ , is a low movable element in soil, therefore its water solubility is a quality index of the fertilizer because only water-soluble phosphorus is absorbed by plant. Fertilizers marked with this icon have a high water-soluble phosphorus content.

---



The use of this range of formulations is characterized by nitrogen protection, reducing losses due to volatilization and leaching (ammonia emissions into the atmosphere). This helps increase the effectiveness of nitrogen fertilization. Fertilizers containing this form of nitrogen are marked with the 'stabilized nitrogen' icon.

---



This type of coated nitrogen allows a targeted supply of fertilizer in the right phenological phase of the crop thanks to a controlled and constant release over time. It improves the effectiveness of use and limits the dosages. Fertilizers containing this form of nitrogen are marked with this icon.

---



The fertilizers marked with this icon contain urea formaldehyde which guarantees a slow release of nitrogen over time, allowing a constant supply of the Nutrient.

---



The organo-mineral fertilizers displaying this icon are obtained starting from organic material containing humified organic substance of high agronomic value.

---



Fertilizers marked with this icon are allowed in organic agriculture.

---




Fertilizers marked with this icon contain microorganisms

---

# LEGEND


# PACKAGING

## Tipology




## Packaging

### Bag




- Bag **1 KG**
- Bag **2,5 KG**
- Bag **5 KG**
  
- Bag **10 KG**
- Bag **20 KG**
- Bag **25 KG**
  
- Big Bag **300 KG**
- Big Bag **500 KG**
- Big Bag **600 KG**

### Tank



- Tank **5 L**
- Tank **20 L**
- Tank **1.000 L**

### Bottle



- Bottle **0,1 L**
- Bottle **0,5 L**
- Bottle **1 L**

# CERTIFICATIONS



## Assofertilizzanti Quality Mark

The Assofertilizzanti Quality Mark is a “collective mark” which allows to identify companies that are committed to operating in quality in the fertilizer production and business system management. It is granted by Assofertilizzanti based on the inspections carried out by the ICQRF, and by MASAF.

Cerea FCP has achieved this mark on all fertilizers produced inside its firm.

---



## Responsible Care

Responsible Care is a voluntary program of the global chemical industry based on the implementation of safety principles and behaviours on Employee health and Environmental Protection and on the commitment to communicate the results achieved, towards a continuous significant and tangible improvement.

Currently, the ‘Responsible Care’ program is adopted by over 10,000 chemical companies in more than 50 countries worldwide, including Cerea FCP

---



## UNI ISO 45001

Cerea FCP has obtained the UNI ISO 45001 certification regarding the occupational health and safety management system. This is the first step to achieve an Integrated Quality, Safety and Environment Management System.

---




## BIOAGRICERT

Cerea FCP has obtained Organic Certification for certain products, issued by Bioagricert, a third-party certification body recognized by the EU. This certification verifies compliance with mandatory regulations and ensures that products and processes meet the required standards for production, processing, and marketing.



# SPECIAL NUTRITION



**NANO.T**

Nano-nutrients

**REACTIVE**

Meso & microelements

**FUTURA**

Biostimulating action

**LEAF**

Macro & mesoelements

# NUTRITIONAL SOLUTIONS

The right products for every need

## Vegetative Development

AIMS

- Lush and balanced growth (vegetative-productive balance)
- Healthy and resistant plants

BEST TO USE

High-quality nutrients, primarily nitrogen, combined with matrices that enhance their efficiency (amino acids, lignosulfonates, etc.)

Leaf N-Fast 

Leaf S-Quality 

Leaf N 

Verv N9  

SuprEmo  

Giove Bio Gold  

## Fruit Size

AIMS

- Uniform size;
- Larger sizes;
- Not compromising quality and shelf life

BEST TO USE

Some particularly pure organic matrices rich in highly biologically active substances (Ashophillum nodosum algae).

Crisco 

VigorGreen  

## Quality of the products

AIMS

- Increase dry matter;
- Improve flavor, texture, and shelf life;
- Intense and uniform fruit color;
- Increase specific weight and protein content

BEST TO USE

Formulations with high-quality calcium and potassium, amino acids and algae, nitrogen and sulfur.

CalcioMagno  

Focus Ca  

Leaf P Ca  

K Fast  

Leaf K 

Leaf S Quality 

Glycos Plus 

VigorGreen  

Nano.T Capo 

MagnetiCal 

Proser Ca  

## Flowering and Fruit Set

AIMS

- Stronger blooms with well-formed flowers;
- Improve fruit set, under stressors
- Reduce fruit drop

BEST TO USE

Elements such as boron and iron and some high-quality organic matrices.

Nanot Fe  

Nanot Fe Bio  

NaturBlack  

Vigorgreen  

B-Power 

StimUp 

Oro-B 

Proser MnZn 

## Rooting

AIMS

To achieve:

- rapid rooting of seeds and successful transplanting;
- healthy, expanded roots;

BEST TO USE

Phosphorus-based formulations and organic matter, essential in this initial phase.

Leaf P Ca  

Hydrostar  

Be Start  

Mycovix  

Verv N9  

SuprEmo   

Nanot Cu 

Nanot Cu Bio  

Giove Bio  

VigorGreen  

## LEGEND

 Organic  
 Foliar application

 Fertigation

## Micronutrient deficiencies

### AIMS

To overcome:

- Nutritional imbalances;
- Specific symptoms due to deficiencies (iron chlorosis, stunted growth, tip burn, etc.);
- Stunted growth

### BEST TO USE

Products containing one or more nutrients, formulated to improve their absorption and translocation.

**Nanot Fe**   
**Nanot Fe Bio**   
**Febo Mix** 

**Febo Bio**   
**Color MgZn**   
**Oro-B** 

**Proser MnZn**   
**Verv Plus** 

## Soil Fertility

### AIMS

To overcome:

- Organic matter deficiency;
- Soil fatigue;
- Compaction

### BEST TO USE

Products based on organic, plant, or animal matrices that improve soil fertility by enhancing the activity of associated products.

**Verv N9**   
**SuprEmo**   
**NaturBlack** 

**Giove Bio Gold**   
**Iride** 

## Salinity

### AIMS

To overcome:

- Limited water absorption capacity by the plant;
- Nutritional imbalances (such as calcium);
- Stunted growth;
- Soil degradation due to high sodium concentrations (sodicity)

### BEST TO USE

Some matrices, such as carboxylic acids, can acidify the soil and sequester some of the salts, including harmful ones like sodium.

**Calcito** 

**Proser Ca** 

## Environmental stress

### AIMS

To overcome: water imbalances; temperature fluctuations; late frosts; phytotoxicity; adverse humidity and temperature conditions.

### BEST TO USE

Products with biostimulant action, containing microorganisms or elements capable of strengthening the plant, improving stress resistance, and promoting crop recovery.

**A-Myco**   
**Mycovix**   
**SuprEmo**   
**NaturBlack** 

**Giove Bio**   
**Cerere**   
**VigorGreen**   
**Proser MnZn** 

**Nanot Cu**   
**Nanot Cu Bio**   
**Nanot Total**   
**Nanot CaPO** 

# TREE CROPS

Nutrient solutions for every phenological phase.

LEGEND

🍷 Organic


FOLIAR

Febo Mix  
Nano.T Cu  
Nano.T Cu Bio 🍷  
Nano.T Total  
A-Myco 🍷


Nano.T Cu (or Nano.T Cu Bio 🍷 ) every 15 days applied every  
Nano.T Total focuses on creating a favorable environment

Giove Bio Gold 🍷  
Leaf N-Fast  
Leaf N


Oro-B 🍷  
NaturBlack 🍷  
Vigorgreen 🍷



VEGETATIVE  
GROWTH



VEGETATIVE  
DEVELOPMENT



PRE FLOWERING  
AND FLOWERING

FERTIGATION

HydroStar  
Verv N9 🍷  
Giove Bio 🍷

SuprEmo 🍷  
Calcito  
Nano.T Fe  
Nano T Fe Bio 🍷

For iron chlorosis Nano.T Fe (or Nano.T Fe Bio 🍷) are the  
Dosage based on the soil's chlorinating power index.

In saline, sodic, calcareous soils with an alkaline pH, add C


every 15 days is advised for stronger and more resilient plants.  
solutions for leaves, shoots and fruits

MagnetiCal  
Crisco  
Leaf P-Ca  
Nano.T CaPO


Leaf K  
Glycos Plus



FRUIT SET



FRUIT SWELLING



VERAISON AND  
RIPENING

CalcioMagno  
Focus Ca

K-Fast

targeted solutions.

Calcito every 20-25 days.

# EXTENSIVE CROPS


Nutrient solutions for every phenological phase.

LEGEND


♥ Organic

FOLIAR

Leaf P-Ca  
SuprEmo ♥  
Leaf N Fast  
NaturBlack ♥  
Giove Bio Gold ♥  
B-Power ♥




FIRST LEAVES




VEGETATIVE  
DEVELOPMENT


Leaf N Fast  
Leaf S Quality  
Leaf N  
Giove Bio Gold 



EARING



FLOWERING



RIPENING



every 15 days is advised for stronger and more resilient plants.  
solutions for leaves, shoots and fruits


Calcio  
Focus Up  
Black  
-B

Giove Bio Gold  
VigorGreen  
MagnetiCal  
Leaf P.Ca


Leaf K  
Proser Ca



FRUIT SET



FRUIT SWELLING



VERAISON AND  
RIPENING

CalcioMagno  
Focus Ca  
Leaf P-Ca

K-Fast  
Giove Bio

targeted solutions.

Calcio every 20-25 days.

# NANO.T TECHNOLOGY FOR AGRICULTURE



Liquid fertilizer family based on nanoparticles in colloidal suspension.

The Nano.T® product line is obtained thanks to the FCP Cerea patented technology. It allows you to correct deficiencies and chlorosis in a new way by improving the efficiency of use of fertilizers by reducing dosages.

# NANO.T®

# NANO.T

Technology for agriculture



CEREA FCP's patented technology allows to produce highly efficient nano-fertilizers.

## 1 High efficiency

Nanoparticles have an elevated contact surface, which eases dissolution and absorption in the plant, allowing a reduced dosage of use;

## 2 Stability in the formulation

The patented production process allows to obtain a stable colloidal suspension over time, avoiding precipitation or aggregation;

## 3 Long-lasting action

Products obtained through the NANO.T process can be employed in the most difficult environmental conditions without altering their characteristics;


## 4 Low environmental impact

Nano-fertilizers do not leach on the soil and have the ability to adhere to the leaf without dispersing in the environment and requiring a limited number of applications;

## 5 Innovative patented technology

The NANO.T technology is an innovation born from a collaboration between the R&D department of FCP Cerea and the department of Biotechnology of the University of Verona.

## Size comparison: from nano to macro



\*1 nanometer = one billionth of a meter (nm = 1m x 10<sup>-9</sup>)


# OPERATING PRINCIPLE

## NANO.T IN FERTIGATION

Nutrients with an internal concentration smaller than 100 mg/kg are called microelements: Iron (Fe), Zinc (Zn), Copper (Cu), Manganese (Mn), Boron (B), Molybdenum (Mo). Four of them (Fe, Zn, Mn, and Cu) oxydise easily into the soil or, reacting with other elements, create insoluble compounds. This causes poor Nutritional efficiency (waste of resources, lower production and environmental impact).

**NANO.T® fertilizers are innovative products that overcome all of these problems.**

The scheme here below shows the operating strategy of NANO.T® fertilizers.



### NANO.T® nutrient release mechanism in the roots

NANO.T® due to its solid nature and its very small dimension (less than 100 nm), doesn't precipitate when it is applied on the soil (which happens with soluble phosphate and sources of Fe, Zn, Mn, Cu) and reaches the roots easily passing through soil pores (it is transported by the mass flow). Once it reaches the rhizosphere (area very close to the plant root), the high ratio surface/volume of NANO.T® helps the solubility - made by roots exudates - and the Nutrients uptake.

## IRON EXAMPLE

Comparison between chelates, complexed and NANO.T® Fe

	NANO.T®	EDDHA o.o.	EDDHA o.p.	EDDHSA	LSA
<b>Photolabile</b> inactivated by the light	NO	YES	YES	YES	NO
<b>Leachable by the soil</b>	NO	YES	YES	YES	NO
<b>Solubility and pH</b>	100% pH 1-10	pH 4- 9 =100%; pH >9= 90 %	pH 4- 9= 90%; pH>10= 80%	pH 4- 9=100%; pH>10 95 %	pH 3.5- 7 =80% pH >8= 50%



### Chelating and complexing agents



Limited persistence since they can be deactivated by the pH in the soil



Reduced effectiveness if exposed to ultraviolet light, heat, oxidizing agents



They can be dispersed in the environment because they are subject to leaching and to polluting groundwater



### NANO.T® Fe



High persistence, effective and stable in a wide range of pH (1-10)



Active in any condition of light and temperature





Low environmental impact as they are retained by the soil and not subject to leaching

# OPERATING PRINCIPLE

## NANO.T WITH FOLIAR APPLICATION

Considering the Nutrients that the plant absorbs in intermediate amounts (meso-nutrients), there are some that the plant struggles to translocate in the most appropriate ways (e.g., calcium). **NANO.T® fertilizers are nanofertilizer formulations that can also provide a solution to this type of problem, thanks to their size and uniformity of distribution.**

The diagram below illustrates the distribution of a NANO.T fertilizer on the leaf.



Mechanism of distribution of NANO.T® on the leaf

NANO.T technology enables each liter of product to contain millions of billions of particles ranging in size from 1 to 100 nanometers, with an extremely high specific surface area. This results in enhanced efficacy and ensures dense leaf coverage (Fig. 2) compared to traditional technology (Fig. 1), at the same application rate.

## THE CALCIUM EXAMPLE

Comparison between calcium chloride and nano-calcium

### Calcium chloride

⊖ It is **poorly mobile within the plant**. It is absorbed by the roots and, through the xylem vessels, tends to settle on the leaves involved in photosynthesis, only reaching a small amount of the fruit through transpiration.

⊖ If applied to leaves, it tends to **stick to their surface**.

### NANO.T® CaPO / Carbo

⊕ **Highly effective at low doses**, nanoparticles have a large contact surface and are evenly distributed across the leaf surface, facilitating absorption.

⊕ **Increasing the calcium content in the rachis** results in a lower percentage of berry detachment in crops such as grapes.


⊕ **Increasing the calcium content in the skin** results in a lower percentage of fruit cracking (e.g., nectarines).

⊕ **Increasing the calcium content in fruit** results in improved quality and shelf life.

## THE COPPER EXAMPLE

CompaRicen with hydroxides, micronized and NANO.T Cu

**Nanoparticles have a dimension one hundred times inferior to that of other products:** in the chart (green curve) the majority of the particles of NANO.T Cu have a dimension of 20 nm while other standard hydroxides (pink curve) measure around 7000 nm as well as the most innovative products such as those micronized (blue curve) around 2000 nm.



### Hydroxides and micronized

- ⊖ They require a **higher dosage** due to their major dimension and inferior uniformity of distribution
- ⊖ **Persistence is reduced** in inverse proportion to the dimension of the particles
- ⊖ They can **disperse in the environment** accumulating in the soil

### NANO.T® Cu

- ⊕ **High effectiveness at low dosages** nanoparticles have a wide surface contact area and distribute uniformly on the leaf, improving absorption
- ⊕ **Persistent on the leaf** the formulation in colloidal suspension allows a better adhesive capacity to foliar waxes
- ⊕ **Lower environmental impact** thanks to the low dosage and the improved adhesive capacity
- ⊕ **It improves the effectiveness of the products combined with it** wide range of miscibility with both fertilizers and/or pesticides



# NANO.T® Fe

## Effectively prevents iron chlorosis

NANO.T® Fe 2.0 effectively prevents iron chlorosis thanks to its formulation containing Nano-Iron in colloidal suspension. NANO.T® Fe 2.0 is persistent because it is retained in the soil (it does not leach) and it is stable in a very wide pH range from 1 to 10. NANO.T® Fe 2.0 stimulates rooting and prevents post-transplant stress. The acidic pH favors the absorption of the fertilizers associated with it. The NANO.T® production process is a patent of FCP Cerea.

### Benefits

- It effectively prevents iron chlorosis, even in soils with high chlorinating power where the chelates are not very effective;
- Promotes an optimal development of the root system;
- Persistent since it is not deactivated (effective at pH 1-10) and is not subject to leaching;
- Easily usable:
  - It can be used during the day as it is not photolabile
  - It does not create deposits in fertigation systems
  - It can be applied both in fertigation and localized with the injector pole
- Acidic pH, increases the effectiveness of the fertilizers associated with it

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Iron (Fe) total</b>	<b>3%</b>
of which soluble in water	2,0%
of which in nano formo	1,0%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>2,0%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>2,5%</b>
<b>Sulfur (SO<sub>2</sub>) soluble in water</b>	<b>3,0%</b>
<b>pH</b>	<b>1,3</b>

### DENSITY

1,16 +/- 0,05 kg/dm<sup>3</sup>

### LOW CHLORINE

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	FERTIGATION DOSAGES	PERIOD
Roots bath	200 ml/hl	pre-transplant
Stone fruits	4-5 l/ha	vegetative growth, pre-flowering, stone swelling, post-harvest.
Actinidia (kiwi)	4-5 l/ha	vegetative growth, pre-flowering, fruit swelling, post-harvest.
Apple tree	4-5 l/ha	vegetative growth, pre-flowering, fruit swelling, post-harvest.
Pear tree	6-10 l/ha	opening of the buds, fruit set, fruit swelling, post-harvest.
Wine and table grapes	6-10 l/ha	vegetative growth, vegetative development, flowering, fruit set.
Citrus trees	30-60 ml/plant	pre-flowering, after fruit set, fruit swelling.
Open field horticultural	3-4 l/ha	2-3 post-transplant applications every 12-15 days.
Tomato	5 l/ha	2-3 post-transplant applications every 12-15 days.
Horticultural in greenhouse	300-500 ml/1000 m <sup>2</sup>	2-3 post-transplant applications every 12-15 days.
All crops	2-3 l/a	FOLIAR APPLICATIONS: any vegetative stage

# NANO.T® Fe Bio

## Effectively prevents iron chlorosis

NANO.T® BIO FE allows a better contribution of iron thanks to nanotechnology. NANO.T® Bio Fe is recommended to prevent iron chlorosis by using an iron control agent. NANO.T® Bio Fe is effective in soils with high chlorinating power characterized by a high content of active limestone. NANO.T® Bio Fe is recommended for the application in fertigation and is also usable in soilless cultivations (peat, cowwithut fiber). The NANO.T® production process is a patent of FCP Cerea.

bioagricert INPUTS



### Benefits

- It effectively prevents iron chlorosis, even in soils with high chlorinating power where the chelates are not very effective;
- Promotes an optimal development of the root system;
- Persistent since it is not deactivated (effective at pH 1-10) and is not subject to leaching;
- Easily usable:
  - It can be used during the day as it is not photolabile
  - It does not create deposits in fertigation systems
  - It can be applied both in fertigation and localized with the injector pole
- Acidic pH, increases the effectiveness of the fertilizers associated with it

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Iron (Fe) soluble in water, complexed with plant extract containing tannins	3%
Sulfur (SO <sub>2</sub> ) soluble in water	5%
pH	2,0

### DENSITY

1,12 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES FERTIGATION	PERIOD
Stone fruits	4-5 l/ha	vegetative growth, pre-flowering, stone swelling, post-harvest.
Actinidia (kiwi)	4-5 l/ha	vegetative growth, pre-flowering, fruit swelling, post-harvest.
Apple tree	4-5 l/ha	vegetative growth, pre-flowering, fruit swelling, post-harvest.
Pear tree	6-10 l/ha	opening of the buds, fruit set, fruit swelling, post-harvest.
Wine and table grapes	6-10 l/ha	vegetative growth, vegetative development, flowering, fruit set.
Citrus trees	30-60 ml/plant	pre-flowering, after fruit set, fruit swelling.
Open field horticultural	3-4 l/ha	2-3 post-transplant applications every 12-15 days.
Horticultural in greenhouse	300-500 ml/1000 m <sup>2</sup>	3-4 post-transplant applications every 15 days.
All crops	2-3 l/a	FOLIAR APPLICATIONS: any vegetative stage



# NANO.T® Cu

## High Efficiency Copper

NANO.T® CU contains copper with high Nutritional efficiency thanks to the small size (nanotechnology) of the particles present in the formulation which makes it effective even at low dosage. NANO.T® Cu 4.5 is ideal for TREATING DEFICIENCIES AND PREVENTING PHYSIOLOGICAL and mechanical DAMAGES (cracks, hail, pruning and harvesting) of the foliar and root system of the plant. NANO.T® Cu 4.5, being a colloidal suspension of nano particles, has a wide contact surface and it does not leach when applied on the leaf and on the soil. NANO.T® Cu 4.5 strengthens the plant and plant tissues.

The NANO.T® production process is a patent of FCP Cera.

### Benefits

- Low-dosage effectiveness thanks to the wide contact surface;
- Low environmental impact as it does not leach if applied on the leaf and in fertigation;
- Low risk of phytotoxicity
- It doesn't stain crops thanks to the innovative formulation;
- High miscibility (fertilizers and / or pesticides), by acidifying the solution.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Copper (Cu) total</b>	<b>4,5%</b>
of which in nano form	2,5%
of which soluble in water	2%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>3,0%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>4,0%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>7,0%</b>
<b>pH</b>	<b>3</b>

### DENSITY

1,18 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

### LOW CHLORINE

## DOSAGES AND USES

CROPS	DOSAGES	MODE
Pome fruit (apple, pear)	1-2 l/ha	foliar
Stone fruit (Peach, Nectarine, Percoco, Apricot, Plum, Almond, Cherry)	1-1,5 l/ha	foliar
Olive (for oil and table). Kiwi, WalNut	3-4 l/ha	foliar
Wine grapes and table grapes	1,5-4 l/ha	foliar
Citrus fruits (Orange, Mandarin, Clementine, Lemon, Grapefruit, Cedar)	3-3,5 l/ha	foliar
HazelNut, ChestNut	3-3,5 l/ha	foliar
Vegetables (processing tomato, table tomato, pepper, aubergine, courgette, cucumber, pumpkin, melon, watermelon, strawberry, artichoke)	3-3,5 l/ha	foliar
Potato, Carrot, Onion, Garlic, Leek, Beetroot	2-3 l/ha	foliar
Spinach, leafy vegetables (lettuce, radicchio, chicory)	2-2,5 l/ha	foliar
Broccoli, Cabbage, Cauliflower, Fennel	1-1,5 l/ha	foliar
Small fruits (blueberry, raspberry, blackberry, etc.)	1,5-2 l/ha	foliar
All crops	2,5-3 l/ha	fertigation

# NANO.T® Cu Bio

## High Efficiency Copper



NANO.T® CU BIO, thanks to the small size (copper hydroxide nano) of the particles present in the formulation is effective even at low dosages. NANO.T® Cu Bio is ideal to treat deficiencies and prevent physiological and mechanical damage (cracks, hail, pruning and harvesting) of the plant's foliar and root system. NANO.T® Cu Bio has a high contact surface and is poorly washable when applied by foliar application and non-washable when applied to the ground. NANO.T® Cu Bio strengthens the plant and plant tissues. The production process NANO.T® is a patent of FCP Cerea.

### Benefits

- Effective at low dosage due to high contact surface;
- Low environmental impact because it is poorly washable when applied by foliar and not leachable when applied in fertigation;
- Low risk of phytotoxicity;
- Does not stain crops thanks to the innovative formulation;
- It improves the effectiveness of the associated products (whether they are fertilisers and/or pesticides) by acidifying the solution.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Copper (Cu) total	5,5%
pH	3,3

### DENSITY

1,10 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES	MODE
Pome fruit (apple, pear)	1-2 l/ha	foliar
Stone fruit (Peach, Nectarine, Percoco, Apricot, Plum, Almond, Cherry)	1-1,5 l/ha	foliar
Olive (for oil and table), Kiwi, Walnut	2-4 l/ha	foliar
Wine grapes and table grapes	2-4 l/ha	foliar
Citrus fruits (Orange, Mandarin, Clementine, Lemon, Grapefruit, Cedar)	3-4 l/ha	foliar
Hazelnut, Chestnut	3-4 l/ha	foliar
Vegetables (processing tomato, table tomato, pepper, aubergine, courgette, cucumber, pumpkin, melon, watermelon, strawberry, artichoke)	2-3 l/ha	foliar
Potato, Carrot, Onion, Garlic, Leek, Beetroot	2-3 l/ha	foliar
Spinach, leafy vegetables (lettuce, radicchio, chicory)	2-2,5 l/ha	foliar
Broccoli, Cabbage, Cauliflower, Fennel	2-2,5 l/ha	foliar
Small fruits (blueberry, raspberry, blackberry, etc.)	1,5-2 l/ha	foliar
All crops	2,5-3 l/ha	fertigation



# NANO.T® Total

## Complete and innovative Nutrition

NANO.T® TOTAL supplies the plant with a complete and innovative Nutrition thanks to the NANO.T production process patented by FCP Cerea. NANO.T® TOTAL promotes root growth, the balanced development of vegetation and, thanks to the presence of iron, it stimulates photosynthesis. NANO.T® TOTAL being in colloidal suspension is neither retrograded nor leached from the ground, resulting effective at low dosages. NANO.T® TOTAL, due to its composition and acidic pH favours root absorption and has an unblocking action on the Nutrients already present in the soil. The NANO.T® production process is a patent of FCP Cerea.

### Benefits

- It helps create favorable environments for maintaining healthy leaves, shoots and fruits;
- It stimulates root development and flowering thanks to the non-retrogradable Nano-phosphorus which is completely available to the plant;
- Improves the absorption efficiency of the root thanks to the acidic pH;
- Promotes vegetative and productive development;
- Increases fruit quality thanks to totally assimilable nitrogen and potassium;
- High environmental effectiveness as it is non-leachable and retrogradable.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>(N) total</b>	<b>6%</b>
of which Nitrogen Ureic	4,4%
of which Nitrogen Nitrico	1,6%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>5%</b>
of which soluble in water	3%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>6%</b>
<b>Iron (Fe) total</b>	<b>2,1%</b>
<b>pH</b>	<b>2,3</b>


### DENSITY

1,21 +/- 0,05 kg/dm<sup>3</sup>

In case of rain, repeat the treatment at the same doses. For foliar treatments, add 1-2 l/ha of NANO.T® Total to the treatment within 7-8 days. NANO.T® Total can be applied at any phenological stage. Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	5 l/ha	during times of greatest need, repeat every 20-30 days
Open field horticultural	5 l/ha	post-transplant, repeat every 15-20 days.
Horticultural in greenhouse	500 ml/ 1000 m <sup>2</sup>	post-transplant, repeat every 15-20 days.



# NANO.T® Carbo

## Fruit quality and shelf life



Liquid fertilizer that allows a high-efficiency calcium supply thanks to nanotechnology, which makes the element readily assimilated by the plant in all its parts and in particular in the fruits. In this way, some physiopathies (apical rot, tip-burn, cracking, etc.) can be prevented and the resistance of plant tissues can be improved, increasing the shelf life of the fruits. In fact, calcium nanoparticles are able to reach the parts of the plant where this element normally struggles to reach. NANO T is a Cerea FCP patent.

### Benefits

- It stimulates root development and flowering thanks to the non-retrogradable Nano-phosphorus which is completely available to the plant;
- Improves the absorption efficiency of the root thanks to the acidic pH;
- Promotes vegetative and productive development;
- Increases fruit quality thanks to totally assimilable nitrogen and potassium;
- High environmental effectiveness as it is non-leachable and retrogradable.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Potassium (K <sub>2</sub> O) soluble in water	9%
Calcium (CaO) total	5,5%
pH	6

Color and density are indicative. For hazard warnings see page 154.

### DENSITY

1,14 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	5 l/ha	during times of greatest need, repeat every 20-30 days.
Open field horticultural	5 l/ha	post-transplant, repeat every 15-20 days.
Horticultural in greenhouse	500 ml/ 1000 m <sup>2</sup>	post-transplant, repeat every 15-20 days.



# NANO.T® CaPO

## Quality and shelf life of fruits

NANO.T® CAPO contains highly efficient calcium, which, thanks to its colloidal nanoparticle formulation, is evenly distributed throughout the plant and is lowly leachable. Calcium is rapidly absorbed and delivered to the fruit, helping to improve its quality and shelf life.

NANO.T® CAPO prevents and treats calcium deficiencies (blossom-end rot, cracking, bitter pit, tipburn) and helps strengthen plant tissue.

### Benefits

- Highly effective at low dosage: thanks to their unique size, the nanoparticles have a large contact surface and are evenly distributed across leaves and fruits, facilitating absorption;
- Persistent on plants and fruits: the colloidal suspension formulation allows for better adhesion to leaf waxes;

It is easy to use as it can be mixed with major plant protection products.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Phosphorus (P <sub>2</sub> O <sub>5</sub> ) total	4%
Potassium (K <sub>2</sub> O) soluble in water	5,5%
Calcium (CaO) total	4,5%
of which soluble in water	2%
pH	3,5

### DENSITY

1,17 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Stone fruits	2-5 l/ha	post-flowering, fruit swelling, veraison
Pome fruits	2-5 l/ha	post fruit set, fruit swelling, veraison
Wine and table grape	3-5 l/ha	pre-flowering, veraison, 15 days before harvest
Kiwi	2-4 l/ha	post fruit set, fruit swelling, 20 days before harvest
Tomato	2-3 l/ha	pre-flowering, post fruit set, veraison
Apple tree, zucchini, cucumber	2-4 l/ha	Post fruit set, fruit swelling
Leafy vegetables	1-3 l/ha	15 days after transplant, 15 days before harvest



# NANO.T® Zn

## Active protection against stress

NANO.T Zn is a liquid formulation based on zinc nanoparticles designed to maximise Nutrient Use Efficiency (NUE). Thanks to its small size (<100 nm), NANO.T Zn overcomes cuticular and stomatal barriers with a much higher translocation rate than conventional zinc. NANO.T Zn is essential for the synthesis of auxin precursors (tryptophan).

It prevents leaf dwarfism and internode shortening.

NANO.T Zn improves the mechanical strength of tissues and the plant's response to abiotic stresses (thermal, water and mechanical).

Thanks to its nanoparticle formulation, NANO.T Zn allows for the controlled release of zinc, ensuring gradual and constant assimilation over time.

### Benefits

- Stimulates the production of secondary metabolites: thicker cell walls (lignification), increased mechanical strength of tissues (plants more resilient to environmental stresses such as temperature changes, drought, wind and/or hail damage);
- Greater photosynthetic efficiency: more vigorous, productive plants and greater leaf expansion;
- Synthesis of auxin precursors: better rooting, flowering and fruit set;
- Greater specific surface area: greater biochemical reactivity, better and more uniform leaf and root coverage, total assimilation without loss due to leaching.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Potassium oxide (K <sub>2</sub> O) soluble in water	5%
Zinc (Zn) total	4%
pH	6

### DENSITY

1,10 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

Crops	Foliar	Fertigation	Period
Fruit trees	2-3 l/ha	5-6 l/ha	Da ripresa vegetativa sino a fine fioritura
Olive trees	2,5-3 l/ha	5 l/ha	Pre-fioritura, fine fioritura, allegagione
Table and grapevine	2-3 l/ha	5-6 l/ha	Germogliamento, pre-fioritura, allungamento grappolo
Open field horticultural	2-2,5 l/ha	4-5 l/ha	Post-trapianto, sviluppo fogliare
Horticultural in greenhouse	2-2,5 l/ha	350-450 ml/1000 m <sup>2</sup>	Post-trapianto, sviluppo fogliare

# NANO.T® Chitosano

## Active defence against stress and drought



NANO.T® Chitosan represents the technological evolution of standard chitosan, a natural polymer derived from chitin. Thanks to its reduction to nanometric scale, its biological and mechanical properties are significantly enhanced.

NANO.T® CHITOSAN is a product with natural elicitor action that increases plant resistance to environmental stress.

NANO.T CHITOSANO also helps plants in case of drought, "helping" them to optimise pore closure during hot hours, reducing water loss.

NANO.T CHITOSANO applied in the pre-harvest phase creates an invisible film that acts as a barrier for the fruit.

### Benefits

- High surface area/volume ratio: greater chemical reactivity and greater ability to penetrate the leaf cuticle, faster and more immediate effect;
- Plants more resistant to environmental stress;
- Healthier fruit before harvesting;
- More productive plants, especially in cases of drought and/or water scarcity.

# NANO.T

**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Chitosan hydrochloride	3%
Degree of deacetylation	> 90 %
Heavy metals	< 40 ppm


### DENSITY

1,02 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative.

## DOSAGES AND USES

Crops	Doses (l/hl) (g/hl of active substance)	Doses (l/ha)	Period	between applications	Mode
Olive tree	0,5 – 1 (15-30)	2-4 l/ha	Da ripresa vegetativa a ingrossamento frutti	2 weeks	Foliar
Table and grapewine	0,5 – 1 (15-30)	2-4 l/ha	Da germogliamento a chiusura grappolo	2 weeks	Foliar
Kiwi	0,5 – 1 (15-30)	2-4 l/ha	Da ripresa vegetativa a ingrossamento frutti	2 weeks	Foliar
Horticultural	0,5 – 1 (15-30)	2-4 l/ha	Da sviluppo vegetativo a ingrossamento frutti	2 weeks	Foliar
Leafy vegetables	0,5 – 1 (15-30)	2-4 l/ha	Da sviluppo foglie a metà ciclo	2 weeks	Foliar
Frutticole (pomacee, drupacee ecc)	0,5 – 1 (15-30)	2-4 l/ha	Da ripresa vegetativa	2 weeks	Foliar
Strawberry and little fruits	0,5 – 2 (15-30)	1-8 l/ha	Da sviluppo vegetativo, a post-allegagione	2 weeks	Foliar
Extensive crops	0,5 – 1 (15-30)	1-4 l/ha	Da sviluppo vegetativo	5-7 days	Foliar



# REACTIVE

## THE MESO & MICROELEMENTS FORCE

Family of formulations with meso and microelements, in liquid and powder form, soluble in water and available for the plants.

The Reactive product line provides plants with the meso- and micronutrients required during the various stages of the growing cycle.

The products in the Reactive line are characterized by meso- and micronutrients complexed with matrices such as carboxylic acids, lignosulfonates, organic acids, and plant matrices.



# Proser MnZn

## Growth promoter

PROSER MnZn is a growth promoter which, by promoting the multiplication of meristematic cells, stimulates the development of both roots and vegetative apices. PROSER MnZn improves the absorption and activity of products applied in combination, amplifying their effectiveness. PROSER MnZn has also an anti-stress action.

### Benefits


- It stimulates vegetative and radical development increasing cell multiplication;
- It promotes fruit development acting on plant metabolism;
- It has an anti-stress action;
- It improves the effectiveness of combined products (fertilizers and/or pesticides), acidifying the solution.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Manganese (Mn) soluble in water	1%
Zinc (Zn) soluble in water	1%
pH	3

### DENSITY

1,08 +/- 0,05 kg/dm<sup>3</sup>

It also contains low molecular weight carboxylic acids.

Do not mix with copper, sulfur, or strongly alkaline reaction products. Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	1-1,5 l/ha	from vegetative growth every 10-15 days	foliar
Wine and table grape	1-1,5 l/ha	from vegetative growth every 10-15 days	foliar
Olive tree	1-1,5 l/ha	from vegetative growth every 10-15 days	foliar
Citrus trees	1-1,5 l/ha	from vegetative growth every 10-15 days	foliar
Open field horticultural	1-1,5 l/ha	from post trapianto every 10-15 days	foliar
Horticultural in greenhouse	90-100 ml/hl	from post trapianto every 5-7 days	foliar
All crops	1-3 l/ha	2-3 applications throughout the cycle	fertigation

# Proser Ca

## Quality and shelf life

bioagricert 



PROSER CA is a specific product for the quality (color and dry substance) and shelf-life of the fruit. The calcium present is withveged and completely assimilated thanks to the action of low molecular weight carboxylic acids. PROSER CA acts as an acidifier, improving the effectiveness of the products combined with it.

### Benefits

- More storable, firm and compact fruits both on the plant and in the post-harvest;
- Increased production, greater dry matter;
- Greater strength of the plant, thanks to the high absorption and transmission of calcium;
- Balanced development of the plant, due to the biostimulating action of the carboxylic acids;
- Improvement of the effectiveness of combined products (fertilizers and/or pesticides), by acidifying the solution.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Calcium (C <sub>3</sub> O) soluble in water	12,5%
pH	1

### DENSITY

1,25 +/- 0,05 kg/dm<sup>3</sup>

It also contains low molecular weight carboxylic acids. Color and density are indicative. For hazard warnings see page 154..

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Stone fruits	2-3 l/ha	from fruit set, every 10-15 days
Pome fruits and Kiwi	2-2,5 l/ha	from after fruit set, every 15-20 days
Wine and table grape	2-3 l/ha	from fruit set, every 15-20 days
Olive tree	2 l/ha	after fruit set, olive swelling, 15 days dalla raccolta
Citrus trees	2-3 l/ha	from fruit set, every 10-15 days
Open field horticultural	2-3 l/ha	da formazione frutticino, every stage interested
Horticultural in greenhouse	100-150 ml/hl	da formazione frutticino, every stage interested
Cereals	2-3 l/ha	with weeds

# Calcito

## High efficiency rhizosphere corrector

CALCITO is a rhizosphere corrector based on short-chain carboxylic acids containing calcium and magnesium. CALCIUM thanks to the long-acting organic acidification of the rhizosphere improves the absorption of Nutrients, in particular calcium, and improves the fertility (structure and pH) of saline, sodic and calcareous soils. CALCITO prevents and treats calcium deficiencies (apical rot, cracking, bitter pit, tip burn) and strengthens plant tissues.

### Benefits


- More balanced and resistant plants (biotic and abiotic stress) thanks to an optimal calcium absorption;
- Better absorption of Nutrients, thank to the long-acting organic acidification of the rhizosphere;
- More capillary rooting and buffering of soil salinity;
- Prevents calcium deficiency diseases: bitter pit, blossom end rot, lettuce hemming;
- More fertile and productive soils as it favors the development of useful microorganisms.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Calcium (CaO) soluble in water</b>	<b>9%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>1%</b>
<b>pH</b>	<b>1</b>

### DENSITY

1,25 +/- 0,05 kg/dm<sup>3</sup>

It also contains low molecular weight carboxylic acids.

Do not mix with products containing phosphorus. Color and density are indicative. For hazard warnings see page 154..

## DOSAGES AND USES

CROPS	DOSAGES FERTIGATION	PERIOD
Stone fruits	10-15 l/ha	preflowering, post fruit set, stone swelling
Pome fruits and kiwi	10-20 l/ha	flowering, after fruit set, fruit swelling
Wine and table grape	10-15 l/ha	vegetative development, fruit set, berry development
Citrus trees	10-15 l/ha	flowering, after fruit set, fruit swelling
Open field fruit vegetables	10-15 l/ha	after 10-15 days from the transplant, repeat every 20 days
Open field Leafy vegetables	10 l/ha	after 7-10 days from the transplant, repeat every 7-10 days
Horticultural in greenhouse	1-2 l/1000 m <sup>2</sup>	after 7-10 days from the transplant, repeat every 7-10 days

# CalcioMagno

## Calcium and magnesium in an acidic formulation for maximum effectiveness

CalcioMagno is a liquid acidic fertilizer (pH 2,6) with a high effectiveness, rich in calcium and magnesium added to organic acids with a carrier action.

CALCIOMAGNO is composed of pure raw materials and is rapidly absorbed and carried within the plant up to the leaves and fruits. Moreover, it provides organic acids that gradually and constantly acidify the rhizosphere improving the assimilation of Nutrients

### Benefits


- Activation of photosynthesis thanks to the presence of magnesium carried and activated by organic acids;
- Improvement of fruit shelf-life thanks to the high withcentration of calcium activated by organic acids;
- Rhizosphere acidification;
- Ideal for horticultural and fruit crops: to tackle apical rot of tomatoes, bitter pit of apples, lettuce hemming and rachis desiccation in the vineyards, and to improve shelf-life and consistency of fruits.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) nitric</b>	<b>8,6%</b>
<b>Calcium (CaO) soluble in water</b>	<b>10%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>4%</b>
<b>pH</b>	<b>2,6</b>

It also contains organic acids. Color and density are indicative. F or hazard warnings see page 154..

### DENSITY

1,41 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	20-30 l/ha	2-3 treatments for stone swelling	fertigation
Pome fruits (apple, pear) and kiwi	20-25 l/ha	2-3 treatments for fruit swelling	fertigation
Wine and table grapes	20-25 l/ha	2-3 treatments for berry swelling	fertigation
Citrus trees	20-30 l/ha	2-3 treatments for fruit swelling	fertigation
Fruit vegetables	20-30 l/ha	2-4 treatments for first stage fruitlets	fertigation
Fruit vegetables	15-20 l/ha	1-3 treatments from mid cycle	fertigation
Greenhouse horticulture	2-3 l/1000 m <sup>2</sup>	1 treatment at fruitlet for each stage	fertigation
All crops	3-5 l/ha	post fruit set	foliar



# Color MgZn

## Greening agent with magnesium sulphur and zinc

Color MgZn is a greening action fertilizer containing magnesium and zinc. Color MgZn is rapidly absorbed, increases the photosynthetic activity and favors the relaxation of plant tissues. Color MgZn has an acidifying action both when used in foliar application and in fertigation. Color MgZn prevents damage due to magnesium deficiency.

### Benefits

- Promotes balanced vegetative development;
- Effectively prevents the onset of rachis desiccation in the vine, phylloptosis on the apple tree, leaf dryness of vegetables;
- It improves the qualitative characteristics thanks to the presence of zinc and sulfur which favor the protein synthesis and aromas;
- It has a rapid effectiveness thanks to the formulation that makes the nutrients easily absorbed and translocated within the plant.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Magnesium (MgO) soluble in water	5%
Sulfur (SO <sub>2</sub> ) soluble in water	10%
Zinc (Zn) soluble in water	1%
pH	5

Color and density are indicative. For hazard warnings see page 154..

### DENSITY

1,21 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees and kiwi	3-4 l/ha	all stages	foliar
Wine and table grape	4-5 l/ha	all stages	foliar
Olive tree	3-4 l/ha	all stages	foliar
Citrus trees	5-6 l/ha	all stages	foliar
Orticole a pieno campo	2-3 l/ha	all stages	foliar
Horticultural in greenhouse	150-200 ml/hl	all stages	foliar
All crops	15-20 l/ha	all stages	fertigation

# Febo Mix

## Meso and microelements activated for photosynthesis

FEBO MIX is a concentrate of meso and microelements complexed with lignosulphonate. FEBO MIX, being rich in magnesium and iron, promotes chlorophyll photosynthesis and has a marked greening activity of the leaves. FEBO MIX activates the plant's metabolism and allows healthy and robust growth.

### Benefits

- Complete and balanced mix for the early stages of plant development (magnesium and iron);
- High effectiveness and assimilability thanks to the presence of lignosulphonates (foliar and root);
- Environmentally friendly since the complexing agents are of vegetal origin with high stability;
- It improves the effectiveness of the combined products (fertilizers and/or pesticides), as, thanks to the lignosulphonates, it has a wetting, tackifying and carrier effect.



**TYPE**  
WATER SOLUBLE  
POWDER



### PACKAGING



### COMPOSITION

Magnesium (MgO) soluble in water	10%
Sulfur (SO <sub>2</sub> ) soluble in water	25%
Boron (B) soluble in water	1,0%
Iron (Fe) soluble in water, complexed with LSA	4,0%
Manganese (Mn) soluble in water, complexed with LSA	1,4%
Molibdeno (Mo)	0,2%
Zinc (Zn) soluble in water, complexed with LSA	1,5%
pH	5,5

### DENSITY

1,00 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	2-3 kg/ha	2-3 applications at leaf emergence and in case of deficiencies	foliar
Pome fruits and kiwi	1-3 kg/ha	2-3 applications at leaf emergence and in case of deficiencies	foliar
Wine and table grape	1,5-2 kg/ha	2-3 applications at vegetative growth and in case of deficiencies	foliar
Olive tree	1-2 kg/ha	2-3 applications at vegetative growth and in case of deficiencies	foliar
Citrus trees	2-3 kg/ha	2-3 applications post-harvest and in case of deficiencies	foliar
Open field horticultural	1,5-2 kg/ha	2-3 applications post-transplant and in case of deficiencies	foliar
Horticultural in greenhouse	150-250 g/hl	2-3 applications post-transplant and in case of deficiencies	foliar
All crops	4-6 kg/ha	2-3 applications	fertigation

# Febo Bio

## Activated microelements for quality productions

bioagricert 



FEBO BIO is a concentrate of trace elements complexed with lignosulphonate. FEBO BIO can be used throughout the production cycle: it promotes optimal plant development, improves nitrogen absorption, thanks to the presence of molybdenum, increases the size, quality and shelf life of the fruit. FEBO BIO is also recommended in post-harvest to restore Nutrient reserves.

### Benefits

- Optimal and balanced development of the plant (iron, manganese and zinc);
- Better nitrogen absorption, thanks to the presence of molybdenum;
- Optimal fruit development, size, quality and shelf life;
- It improves the effectiveness of the combined products (fertilizers and/or pesticides), as, thanks to the lignosulphonates, it has a wetting, tackifying and carrier effect.



**TYPE**  
WATER SOLUBLE  
POWDER



### PACKAGING



### COMPOSITION

Boron (B) soluble in water	0,7%
Iron (Fe) soluble in water, complexed with LSA	4,0%
Manganese (Mn) soluble in water, complexed with LSA	3,5%
Molibdeno (Mo)	0,2%
Zinc (Zn) soluble in water, complexed with LSA	2,0%
pH	5

### DENSITY

1,00 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154..

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	2-2,5 kg/ha	2-3 applications throughout the cycle	foliar
Pome fruits and kiwi	2-2,5 kg/ha	2-3 applications throughout the cycle	foliar
Wine and table grape	1,5-2 kg/ha	2-3 applications throughout the cycle	foliar
Olive tree	1,5-2 kg/ha	2-3 applications throughout the cycle	foliar
Citrus trees	2-2,5 kg/ha	2-3 applications throughout the cycle	foliar
Open field horticultural	2-2,5 kg/ha	2-3 applications throughout the cycle	foliar
Horticultural in greenhouse	90-100 g/hl	3-4 applications throughout the cycle	foliar
All crops	4-5 kg/ha	3-4 applications throughout the cycle	fertigation

# MagnetiCal

## Foliar energizer based on calcium and magnesium

MagnetiCal is a liquid fertilizer rich in sugar which readily provides energy to the plant.

The sugar and the subacidic pH favour a rapid and complete assimilation of calcium and Magnesium. The use of MagnetiCal increases the vigourness of the plant (better photosynthesis and metabolic activity), by strengthening and increasing the consistency of fruits; in this way it is possible to obtain a product of much higher quality.

### Benefits

- The plant is more active thanks to the significant presence of sugar and magnesium;
- The plant is more robust and fruits have a better shelf-life thanks to the presence of calcium;
- Organoleptic qualities are enhanced (flavour, consistency and colour);
- It improves the effectiveness of other products employed (both fertilizers and/or pesticides), since it has an adhesive and carrier capacity.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Calcium (C <sub>3</sub> O) soluble in water	12%
Magnesium (MgO)	4%
pH	5

### DENSITY

1,30 +/- 0,05 kg/dm<sup>3</sup>

It also contains plant mono and polysaccharides. Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	3-4 l/ha	3-4 applications from fruit set every 15 days	foliar
Pome fruits (Apple tree, Pear tree) e kiwi	3-4 l/ha	3-4 applications from fruit set every 15 days	foliar
Wine and table grape	3-4 l/ha	3-4 applications from fruit set every 15 days	foliar
Citrus trees	3-4 l/ha	3-4 applications from fruit set every 15 days	foliar
Olive tree	3-4 l/ha	3-4 applications from fruit set every 15 days	foliar
Open field fruit vegetables	3-3,5 l/ha	3-4 applications from fruit set in every stage	foliar
Open field Leafy vegetables	3 l/ha	2-3 applications from 5-6 leaves every 8-10 days	foliar
Horticultural in greenhouse	250-300 ml/hl	3-4 applications from fruit set every 7 days	foliar
All crops	10 - 20 l/ha	2-3 applications	fertigation

# Focus Ca

## Calcium organo-complessato sistemico

FOCUS Ca contains a high percentage of calcium totally complexed with ligninsulfonate; this makes the calcium completely assimilable, mobile inside the plant and not washable in the soil. The product is effective at low doses even at low temperatures and, thanks to its acidic pH, it can improve the absorption of Nutrients in the soil.

FOCUS Ca is particularly suitable to combat the apical rot, bitter buteratura of pome fruits, tip burn of salads and helps to counteract the desiccation of the rachis in vineyards. FOCUS Ca directly affects the quality of the productions improving the shelf life both on the plant and in the post-harvest.



**TYPE**  
Liquid



### PACKAGING



### Benefits

- Stronger and more robust plants (thicker and more resistant plant tissues);
- Better fruit quality (colour, taste);
- Increased shelf life on the plant and in post-harvest;
- Reduced susceptibility to fruit cracking.

### COMPOSITION

<b>Calcium (CaO) soluble in water</b>	<b>15%</b>
Complexed with ammonium lignosulfonate	12%
<b>pH</b>	<b>1</b>

### DENSITY

1,45 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154..

## DOSAGES AND USES

CROPS	DOSAGES RADICALI	DOSAGES FOGLIARI	PERIOD
Pome fruits	20-25 l/ha	3-4 l/ha	From fruit set every 15 days
Stone fruits	20-25 l/ha	3-4 l/ha	From fruit set every 10-12 days
Olive tree	20-25 l/ha	3-5 l/ha	From fruit set every 15-20 days
Wine and table grape	15-20 l/ha	3-4 l/ha	From fruit set every 15 days
Citrus trees	15-20 l/ha	3-4 l/ha	From fruit set every 15-20 days
Actinidia (kiwi)	20-25 l/ha	3-5 l/ha	From fruit set every 15 days
WalNut, hazelNut, chestNut	15-20 l/ha	3-4 l/ha	From fruit set every 15 days
Open field horticultural	15-20 l/ha	3-4 l/ha	From fruit set to first stage every 8-10 days
Horticultural in greenhouse	1-2 l/1000 m	200-250 ml/100 l of waters	From fruit set to first stage every 8-10 days
Open field Leafy vegetables	10-15 l/ha	2,5-3 l/ha	From post-transplant to 8-10 days after harvest
Potato, carrot, onion, garlic, leek	10-15 l/ha	3-4 l/ha	From tuber formation every 15 days



# Oro-B

## Complexed Boron for flowering and fruit set

ORO-B is a liquid fertilizer composed of Boron complexed with ethanolamine (an organic molecule that stabilizes and facilitates absorption). This formulation ensures rapid and uniform delivery of Boron within plant tissues. Boron is an essential microelement, particularly important in the flowering and fruit set stages and contributes to cell wall formation and tissue lignification. Furthermore, Boron actively intervenes in several biochemical processes including nucleic acid synthesis and carbohydrate metabolism.

### Benefits

Stimulates and evens out flowering:

- Improves and promotes fruit set
- Improves sugar content and fruit quality (translocation of sugars and calcium)
- Makes the stems and petioles more robust and resistant to mechanical (wind) and environmental stresses.



**TYPE**  
Liquid



### PACKAGING



TRANSPARENT



### COMPOSITION

Boron (B), ethanolamine, water soluble	11%
pH	8

Color and density are indicative. For hazard warnings see page 154.


### DENSITY

1,37 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	1-2 l/ha	pre-flowering / Post-harvest	Foliar
Pome fruits	1-2 l/ha	pre-flowering / Post-harvest	Foliar
Wine and table grape	1-2 l/ha	pre-flowering / Post-harvest	Foliar
Citrus trees	1-2 l/ha	pre-flowering / Post-harvest	Foliar
Nut, hazelnut	1-2 l/ha	pre-flowering / Post-harvest	Foliar
Olive tree	1-2 l/ha	Pre-flowering, flowering, end of flowering, post-harvest	Foliar
Open field fruit vegetables	1-2 l/ha	Flowering	Foliar
Horticultural in greenhouse	1-2 l/ha	Flowering	Foliar
Extensive crops	1-2 l/ha	Pre-flowering	Foliar
All crops	4-5 l/ha	pre-flowering 2-3 interventions depending on the severity of the deficiency	fertigation, only in case of severe Boron deficiency in the soil





# FUTURA

## BIOSTIMULATING ACTION BY NATURE

Family of liquid and powder products containing active organic substances which stimulate the physiological process of the plant, promoting growth and productivity.

The Futura family includes products based on amino acids, algae extracts, and humic acids, whose function is to stimulate the natural processes of plants to increase nutrient absorption, tolerance to abiotic stress, and crop quality.

# Verv Plus

## Biopromoter of vegetative development

Verv Plus is a bio-promoter of vegetative development containing over 45% of total amino acids in mainly levorotatory form. Verv Plus activates the plant's metabolism thanks to the presence of organic nitrogen of amino acid origin and a balanced mix of microelements. Verv Plus improves productivity by stimulating vegetative development, flowering and fruit set.

### Benefits

- Improves productivity by stimulating vegetative development, flowering and fruit set;
- It has a biostimulating action thanks to the presence of free amino acids and oligopeptides;
- It activates the metabolism of the plant thanks to the mix of microelements, preventing the onset of deficiencies;
- It increases the effectiveness of the products distributed together, thanks to the co-formulating action of the amino acid matrix.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Nitrogen (N) organic	7%
Carbon (C) organic	22%
Boron (B) soluble in water	0,5%
Iron (Fe) soluble in water	1%
Zinc (Zn) soluble in water	0,5%
pH	7,5

Color and density are indicative. For hazard warnings see page 154.

### DENSITY

1,25 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	1,5-2,5 l/ha	2-3 applications every 10-20 days from vegetative growth
Wine and table grape	1,5-2,5 l/ha	2-3 applications every 10-20 days from vegetative growth
Horticultural	1-2 l/ha	3-4 applications starting 10-15 days
Extensive crops	1,5-2,5 l/ha	after transplanting



# VerV N9

## Promoter of radical activity


VERV N9 is an organic nitrogen fertilizer with a very high content of amino acids, oligopeptides and peptides, greater than 50%, obtained from enzymatic hydrolysis (low salinity). VERV N9 has a phytostimulant action and a very high Nutritional efficiency, promoting the development and activity of both roots and the foliar apparatus. VERV N9 improves soil fertility conditions and stimulates the growth of microorganisms and their activity.



**TYPE**  
Liquid



### PACKAGING



Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	10-20 l/ha	vegetative growth, in pre-flowering, post fruit set, fruit development	fertigation
Open field horticultural	10-30 l/ha	Post-transplant post fruit set, fruit development	fertigation
Horticultural in greenhouse	2-4 l/1000m <sup>2</sup>	3-4 applications after transplant every 10-15 days	fertigation
Cereals	3-4,5 l/ha	Tillering, flag leaf	foliar
All crops	2,5-5 l/ha	vegetative development	foliar

bioagricert <sup>INPUTS</sup> ✓



### Benefits

- Stimulating effect thanks to the high content of amino acids, oligopeptides and peptides > 50%;
- Promotes the development of the plant in particular of the roots;
- It increases the efficiency of the fertilizers used in combination;
- More constant release and limited risk of leaching;
- It increases the biological fertility of the soil thanks to the quality of the organic matrix (low salinity);
- Versatility since it can be used both in fertigation and foliar application in association with other products for Nutrition and defence.

### COMPOSITION

<b>Nitrogen (N) organic</b>	<b>9%</b>
<b>Carbon (C) organic</b>	<b>24,5%</b>

### ALSO CONTAINS:

total amino acids	50%
of which Glycine	13%
of which Proline	7,5%
of which Hydroxyproline	6%
of which Glutamic Acid	5%
of which Arginine	5%

<b>pH</b>	<b>5,5</b>
-----------	------------

### DENSITY

1,25 +/- 0,05 kg/dm<sup>3</sup>

# Glycos Plus

## Promoter of coloring and brix degree

Glycos plus is a specific product to promote and uniform the color and brix degree of fruits. Glycos plus has a formulation based on amino acids and potassium designed to induce fruit ripening while maintaining its shelf life. Glycos plus also contains boron which promotes the absorption of potassium and amplifies its effectiveness.

### Benefits

- Promotes and uniform fruit color, increases quality and optimizes harvesting;
- It increases the brix degree of the fruit (sugar content) improving its shelf life;
- Maintains shelf life as it does not overripen the fruit;
- Easy to use, it can be applied mixed with all the main foliar products



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>5%</b>
of which organic	3%
of which ureic	2%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>9%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>14%</b>
<b>Carbon (C) organic</b>	<b>8%</b>
<b>Boron (B) soluble in water</b>	<b>0,25%</b>
<b>pH</b>	<b>8</b>

Color and density are indicative

### DENSITY

1,25 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Peach, apricot, Apple tree	2 l/ha	45, 30 and 15 days before the harvest
Wine and table grape	2 l/ha	40% berry development, beginning of veraison, after 10 days
Strawberry	2 l/ha	blossoming, green fruit and veraison from white to red

# StimUp

## Growth bioactivator



StimUp is a specific action product with an activating action. StimUp stimulates vegetative and root development, blossoming and fruit set thanks to the hormone-like activity. StimUp is effective at low doses as it contains short-chain humic and fulvic acids obtained by distillation that are more active and easily absorbed by the plant.

### Benefits

- Stimulates vegetative and root development thanks to hormone-like activity;
- Increases flowering and fruit setting thanks to the biostimulant action;
- High effectiveness at low dosages as it contains short-chain humic and fulvic acids obtained by distillation that are more easily absorbed by the plant;
- Versatility, it can be combined with all the main liquid products on the market (fertilizers and pesticides).



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Organic matter on fresh weight	2,2%
Organic matter on dry weight	61%
Percentage of humified substance on organic matter	97%
Organic nitrogen on dry weight	0,5%
C/N Ratio	53%
pH	6,5

Color and density are indicative

### DENSITY

1,01 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	120-200 ml/ha	at the beginning of blossoming. if necessary, a seewithd application in full bloom
Citrus trees	150-200 ml/ha	at the beginning of blossoming. if necessary, a seewithd application in full bloom
Wine and table grape	120-170 ml/ha	at the beginning of blossoming. if necessary, a seewithd application in full bloom
Olive tree	150 ml/ha	end of blossoming
Tomato	200-250 ml/ha	stress conditions
Open field horticultural and in greenhouse	20-25 ml/hl	stress conditions
Potato	100 ml/ha	foliar development
Seed treatment	5 ml/kg	before the seeding
All crops	100 ml/hl	roots bath

# Cerere

## Thermoprotector and osmoregulator



Cerere acts as a thermoprotector and osmoregulator, it is composed of short-chain polyols and humic extracts of natural origin. Cerere is indicated to limit damage from abiotic stress, in particular cold, heat, excessive salinity, root asphyxia and hail (after the event occurs, it helps the plant to recover faster).

### Benefits

- Increases resistance and reduces recovery time in the event of damage from cold, heat, excessive salinity, root asphyxiation and hail;
- It safeguards quantity and quality even in case of too low or too high temperatures, excesses of light and salinity stress;
- Limits damage from cracking and burning;
- Saving of water, possibility of lengthening the irrigation intervals as it increases the resistance of the crop to water stress;
- Effective on all crops;
- Easy to use, it can be applied mixed with all the main products.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

47% of medium and long polymer chain polyols	47%
pH	6,5

Color and density are indicative. For hazard warnings see page 154.

### DENSITY

1,10 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	1,5-2,5 l/ha	1 application 24-48 hours before the stressful event, then every 7-10 days
Wine and table grape	1,5-2,5 l/ha	1 application 24-48 hours before the stressful event, then every 7-10 days
Open field horticultural and in greenhouse	1-2,5 l/ha	1 application 24-48 hours before the stressful event, then every 7-10 days

# B-Power

## Bioactivator for flowering and fruit set



B-POWER is a bioactivator that thanks to the particular formulation based on natural components of plant origin performs a double action: it activates the plant metabolism and has anti-stress effect. B-POWER contains Boron complexed with humic and fulvic acids, thus being totally assimilable and withveyed inside the plant. B-POWER stimulates radical development, increasing Nutrient absorption and promoting flowering and fruit sets.

### Benefits

- Activates the plant, promoting flowering and fruit sets;
- Stimulates radical development;
- Increases Nutrient absorption;
- It performs anti-stress action.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Boron (B) soluble in water</b>	<b>2%</b>
<b>humic and fulvic acids estratti with acqua</b>	<b>14%</b>
<b>pH</b>	<b>7,5</b>

Color and density are indicative. For hazard warnings see page 154.

### DENSITY

1,10 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	3-4 l/ha	opening of the buds fiorali, flowering, post fruit set, in the process of swelling the fruit	foliar
Pome fruits and kiwi	2-3 l/ha	pre-flowering, flowering, post fruit set, in the process of swelling the fruit	foliar
Grapewine and table grape	3 l/ha	pre-flowering, flowering, grano di pepe, in the process of berry enlargement	foliar
Olive tree	4 l/ha	pre-flowering, flowering, cluster flowers, olive swelling phase	foliar
Citrus trees	2-2,5 l/ha	pre-flowering, flowering, post fruit set, in the process of swelling the fruit	foliar
Open field horticultural	2-2,5 l/ha	before and after la flowering, in the process of fruit swelling	foliar
Horticultural in greenhouse	300-500 ml/1000 m <sup>2</sup>	before and after la flowering, in the process of fruit swelling	foliar
HazelNut tree	3-4 l/ha	start of leaf fall	foliar
Extensive crops	3-6 l/ha	with pesticides	foliar
All crops	4-6 l/ha	before and after la flowering, in the process of fruit swelling	FERTIGATION

# K-Fast

## Potassium activated for ripening



K-FAST contains a high content of activated potassium. K-FAST thanks to the synergistic action of sulfur and humic and fulvic acids, acidifies the rhizosphere allowing complete absorption of potassium which is rapidly translocated in the fruit. K-FAST also performs an anti-stress action supporting the plant in the delicate phase of fruit ripening, allowing to harvest quality fruit with a higher dry matter content, limiting the risk of over-ripening.

### Benefits

- Stimulates ripening, uniforms and increases color and brix degree thanks to the high content of soluble and rapidly absorbable potassium;
- High efficiency since the potassium complexed by the organic matrix limits the risk of leaching;
- It has a bio-stimulating and stress-protective action thanks to the presence of humic and fulvic acids;
- Increases Nutrition efficiency thanks to a better root development and an acidic pH.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Potassium (K <sub>2</sub> O) soluble in water	24%
Sulfur (SO <sub>3</sub> ) soluble in water	40%
Contains humic acids	
pH	8,5

Color and density are indicative

### DENSITY

1,45 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits and pome fruits	10-30 l/ha	3-4 applications from ripening	fertigation
Wine and table grape, hazelNut	10-20 l/ha	3-4 applications from veraison	fertigation
Open field horticultural	10-20 l/ha	3-4 applications from ripening	fertigation
Horticultural in greenhouse	1-3 l/1000m <sup>2</sup>	Every 5-7 days from the start of ripening	fertigation
All crops	3-5 l/ha	3-4 applications from ripening	foliar

# Edafos

## Betainphosphate from roots to fruit

EDAFOS contains a high title of phosphorus complexed with betaines. EDAFOS has a starter action, promotes rooting, stimulates blossoming and strengthens the plant. The combined action of phosphorus and betaines makes the plant less susceptible to stressful conditions. EDAFOS thanks to the very acidic pH creates an optimal environment for the absorption of nutrients. If used on stone fruits after fruit setting, it stimulates the growth and the hardening of the stone. On the vine, in post-fruit set, it stimulates the elongation of the bunch.

### Benefits


- Better root development, increase and synchronization of blossoming, better fruit set thanks to the high content of soluble and rapidly absorbable phosphorus;
- High efficiency as the phosphorus complexed by the organic matrix is not retrograded and leaching is limited;
- Biostimulating action and protection from stress thanks to the presence of betaines;
- Better nutrition efficiency thanks to better root development and acidic pH.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

**Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water** 40%

**With betaine**

**pH** 1

Color and density are indicative

### DENSITY

1,42 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	10-25 l/ha	alla vegetative growth, in pre-flowering, post fruit set, fruit swelling
Open field horticultural	10-30 l/ha	post-transplant, pre-flowering, fruit set
Horticultural in greenhouse	2-3 l/1000m <sup>2</sup>	3-4 3-4 applications after the transplant each 10-15 days

# Be-Start 5.15

## High efficiency liquid starter with betaines

BE START 5.15 is a ready-to-assimilate liquid starter. BE START 5.15 is characterized by a balanced content of nitrogen and phosphorus. The phosphorus being bound to the betaines, is conveyed more efficiently and is not leached reducing the risk of retrogradation. BE START 5.15 stimulates rooting, strengthens the plant, promotes flowering and fruit setting and, if applied during fruit development, it promotes enlargement and hardening of the stone; in the vineyards, it favors the lengthening of the bunch and strengthens the petiole.

### Benefits

- High performance liquid starter, readily assimilable nitrogen and phosphorus since they are totally soluble;
- High efficiency since nitrogen and phosphorus complexed by the organic matrix are not retrogradable and not leachable;
- Biostimulating action and protection from stress thanks to the presence of betaines;
- Versatility, as it can be applied both by foliar and root application.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) toale</b>	<b>5%</b>
of which ammoniacal	3%
of which ureic	2%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>15%</b>
<b>Contiene betaine</b>	
<b>pH</b>	<b>4</b>

Color and density are indicative

### DENSITY

1,15 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD	MODE
Fruit trees	10-20 l/ha	at vegetative growth, pre-blossoming, post fruit set, fruit swelling	fertigation
Open field horticultural	10-15 l/ha	post-transplant, post fruit set, fruit swelling	fertigation
Horticultural in greenhouse	2-3 l/1000m <sup>2</sup>	3-4 applications from 10-15 days after transplant	fertigation
Cereals	3-5 l/ha	post-emergency, tillering	foliar
All crops	3-6 l/ha	2-4 applications according to the needs of the crop	foliar

# Iride

## Energy promptly available

IRIDE contains organic phosphorus in the form of vegetal origin, with a stimulating action. IRIDE, providing phosphorus readily assimilable by the plant and humic and fulvic acids (24%), facilitates rooting and vegetative development. Being rich in carbohydrates, IRIDE gives energy to both the plant and the soil, promoting the development of useful microorganisms and increasing fertility.

### Benefits



**TYPE**  
Liquid



### PACKAGING



Contains humic and fulvic acids. Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	10-20 l/ha	vegetative growth, pre-blossoming, after fruit set, fruit swelling	fertigation
Open field horticultural	10-30 l/ha	after transplant, after fruit set, fruit swelling	fertigation
Horticultural in greenhouse	2-4 l/1000m <sup>2</sup>	3-4 applications after transplant each 10-15 days	fertigation
Cereals	3-5 l/ha	tillering, flag leaf	foliar
All crops	3-6 l/ha	2-4 applications according to the needs of the crop	foliar

bioagricert



- Energy booster stimulates root development, flowering and vegetative development;
- Biostimulating action thanks to the combined action of a high content of humic and fulvic acids 23% and betaines;
- Rapid effectiveness since nitrogen, phosphorus and potassium, being of organic origin, are totally soluble and readily assimilable;
- High efficiency since nutrients in organic form are difficult to retrograde and leach

### COMPOSITION

Nitrogen (N) organic	3%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) total	4%
Potassium (K <sub>2</sub> O) soluble in water	4%
Carbon (C) organic	14,5%
pH	4,5

### DENSITY

1,20 +/- 0,05 kg/dm<sup>3</sup>

# Giove Bio N8

## 100% vegetal physioactivator

bioagricert INPUTS ✓



GIOVE BIO is a 100% vegetal concentrated physioactivator rich in vegetal amino acids (about 98%), obtained by enzymatic hydrolysis. GIOVE BIO thanks to the high content of organic nitrogen, readily assimilable, promotes a balanced vegetative development. GIOVE BIO is rich in glycine, an amino acid with a carrier action, which improves the effectiveness of the products associated with it, and proline, an amino acid with an anti-stress action, which promotes growth and productivity even in the event of unfavorable climatic conditions.

### Benefits

- It has a biostimulating action thanks to the high presence of peptides, oligopeptides and free amino acids, all of vegetal origin;
- It activates the metabolism of the plant thanks to the high content of totally soluble organic nitrogen;
- It's environmentally friendly and innovative as it comes from the enzymatic hydrolysis of plant tissues;
- It increases the effectiveness of the products distributed together thanks to the co-formulating action of the amino acid matrix.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) organic</b>	<b>8%</b>
<b>Carbon (C) organic</b>	<b>25%</b>
<b>pH</b>	<b>6,5</b>

### ALSO CONTAINS:

total amino acids	50%
-------------------	-----

Color and density are indicative

### DENSITY

1,20 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	2-3 l/ha	2-3 applications from vegetative development	foliar
Wine and table grape	2-3 l/ha	2-3 applications from vegetative development	foliar
Olive tree	3-4 l/ha	vegetative growth, fruit swelling	foliar
Open field horticultural	2-3 l/ha	every 7 -10 days from transplant	foliar
Horticultural in greenhouse	250-300 ml/hl	every 5-7 days from transplant	foliar
Cereals	2-3 l/ha	booting, flag leaf	foliar
Fruit trees	10-15 l/ha	from vegetative growth	fertigation
Open field horticultural	10-12 l/ha	from post-transplant	fertigation
Horticultural in greenhouse	1-1,5 l/1000 m <sup>2</sup>	from post-transplant	fertigation

# NaturBlack

## Highly concentrated activator of humic acid extracts

bioagricert 



NATURBLACK is a concentrate of humic extracts (28% of which 22% humic and 6% fulvic) with an activating action on the plant. NATURBLACK applied on the leaves has an auxin-like physioactivating action that is particularly strong in situations of high stress for the plant (weeding, sudden changes in temperature, pruning, hail, etc). NATURBLACK used in fertigation stimulates rooting, counteracts post-transplant stress of seedlings, promotes the absorption of nutrients blocked in the soil, improves soil structure and it is an ideal substrate for the proliferation of useful soil microorganisms. NATURBLACK, having an acidic pH, is optimal both in combination with pesticides and other fertilizers.

### Benefits

- Stimulates the growth and development of the plant by activating its metabolism;
- It has an anti-stress effect for the plant (transplant, weeding, sudden changes in temperature, pruning, hail, frost);
- Promotes rooting and facilitates the absorption of nutrients present in the soil;
- Improves the physical fertility of the soil (structure) and the biological fertility by promoting the development of useful microorganisms in the soil;
- Improves the effectiveness of the products combined with it (fertilizers, pesticides) thanks to the acidic pH.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Organic carbon on dry weight	30%
Organic Carbon extractable on total organic Carbon	60%
Organic humified Carbon on extractable organic Carbon	60%
Organic nitrogen on dry matter	0,5%
Organic matter on dry matter	90%
Organic matter extractable on % of the organic matter	60%
Organic matter humified on % of the extractable matter	60%
pH	3

Extracting medium water. Color and density are indicative

### DENSITY

1,10 +/- 0,05 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
All Crops	1-2 l/ha	pre-transplant	bagnetto
Fruit trees	1-2 l/ha	from vegetative growth	foliar
Wine and table grape	1-1,5 l/ha	every 12-15 days	foliar
Olive tree	1-2 l/ha	from vegetative growth	foliar
Open field horticultural	1-1,5 l/ha	every 12-15 days	foliar
Horticultural in greenhouse	100-150 ml/hl	from vegetative growth	foliar
Cereals	1-2 l/ha	every 12-15 days	foliar
All crops	3-5 l/ha	from transplant every 8-10 days	fertigation

# Giove Bio Gold

## 100% vegetable amino acids

bioagricert 



GIOVE BIO GOLD has a very high concentration of vegetal amino acids (98%) in medium-short chains with a phytostimulant, anti-stress and nutritional action. GIOVE BIO GOLD is effective at low doses thanks to the highly soluble powder formulation. GIOVE BIO GOLD is rapidly absorbed and activates the plant's metabolism allowing for optimal development and rapid recovery of stressed and/or blocked plants.

### Benefits

- High effectiveness at low dosages thanks to the very high concentration;
- Biostimulating action thanks to the high presence of peptides, oligopeptides and free amino acids, all of vegetal origin;
- Activation of plant metabolism thanks to the high content of totally soluble organic nitrogen;
- Environmentally friendly and innovative as it comes from the enzymatic hydrolysis of plant tissues;
- Easy to use thanks to the high solubility;
- Increased effectiveness of the products distributed together thanks to the co-formulating action of the amino acid matrix.



### TYPE

Water-soluble powder

### PACKAGING



### COMPOSITION

<b>Nitrogen (N) organic</b>	<b>16%</b>
<b>Carbon (C) organic</b>	<b>50%</b>
<b>pH</b>	<b>6,5</b>

### ALSO CONTAINS:

<b>Total amino acids</b>	<b>98%</b>
of which Glicina	25,5%
of which Prolina	14%

Color and density are indicative

### DENSITY

0,60 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	1-2 kg/ha	vegetative growth, before blossoming, after fruit set, fruit development	foliar
Wine and table grape	1-2k g/ha	pre-blossoming, blossoming, berries goat-sized, bunches begin to hang, berries swelling	foliar
Olive tree	1-2 kg/ha	pre-blossoming, blossoming, cluster flowers, drupe swelling	foliar
HazelNut	2 kg/ha	from fruit set	foliar
Open field horticultural (Potato, Tomato, carciofo...)	1-3 kg/ha	post-transplant/ post emergence, vegetative development, post fruit set, fruit development	foliar
Horticultural in greenhouse	300-500 g/1000m <sup>2</sup>	3-4applications 10-15 days after transplant	foliar
Cereals	1-2 kg/ha	tillering, leaf flag	foliar
Potato	1 kg/ha	10-15 days after emergence until tuberification	foliar
All crops	5-6 kg/ha	2-4 applications	fertigation

# SuprEmo

## Plant and soil physioactivator



SUPREMO is a product containing pure hemoglobin deriving from treated and concentrated animal blood. The low-temperature processing allows to preserve all functional properties of hemoglobin by ensuring an optimal effectiveness on the field. SUPREMO is rich in essential amino acids and in a significant quantity of tryptophan, an amino acid precursor of auxins, stimulating the plant development.

SUPREMO is particularly rich in organic iron, which is very active biologically and its constant use allows to maintain plants green and active.

SUPREMO can be used in any types of soil (alkaline, saline, sodic, etc.). It stimulates the activity of useful

microorganisms already present in the soil or inoculated in it, which help release the nutrients. SUPREMO may be employed with low temperatures and is specifically suitable for stimulating plants "blocked" in their development.

### Benefits

- Rapid development of plants thanks to the prompt action, even at low temperatures;
- Increases fruit quality improving color and swelling;
- Increases the microbiological fertility of the soil releasing nutrients;
- Versatile use, suitable both in foliar applications and fertigation.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) organic</b>	<b>5%</b>
<b>Carbon (C) organic</b>	<b>18%</b>
<b>Iron (Fe) organic</b>	<b>800 mg/kg</b>
<b>pH</b>	<b>5,5</b>

### DENSITY

1,10+/- 0,05 kg/dm<sup>3</sup>

Preservative used: trisodium citrate.. Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	20-40 l/ha	from vegetative growth	fertigation
Pome fruits	20-40 l/ha	from vegetative growth	fertigation
Kiwi	20-40 l/ha	from vegetative growth	fertigation
Olive tree	20-40 l/ha	from vegetative growth	fertigation
Wine and table grape	20-40 l/ha	from vegetative growth	fertigation
Horticultural in greenhouse	2-3 l/1000 metri	from post-transplant	fertigation
Open field horticultural	20-30 l/ha	from post-transplant	fertigation
Cereals	5 l/ha	with weeding	foliar
All crops	3-5 l/ha	everytime	foliar

# Crisco

## Initiator of cell division

CRISCO is a concentrate of Canadian *Ascophyllum Nodosum* algae (25%), rich in biologically active substances that promote cell division. Thanks to its composition, it activates photosynthesis increasing the content of chlorophyll in the leaves.

Cell division has a direct effect on the fruit, since it contributes to a greater size, and an indirect effect because it contributes to swelling and rachis elongation with a consequent easier penetration of nutrients.

Moreover, algae extracts have a biostimulant effect on the plant. Increasing metabolism and their capacity of facing more effectively potential stress conditions.

### Benefits

- Increase yields thanks to a major size;
- Stimulates plant activity even in unfavourable conditions;
- Safeguards the health of perennial plants thanks to the natural hormone-like action.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Carbon (C) organic	12%
Mannitol	7,5 g/L
pH	7

### DENSITY

1,20 +/- 0,05 kg/dm<sup>3</sup>

it contains *Ascophyllum Nodosum* Canadese algae (25%). Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Apple tree e Pear tree	1 l/ha	increasing fruit size: post-fruit setting, after physiological drop
Citrus trees, Olive tree, Stone fruits	1 l/ha	increasing fruit size: post-fruit setting, fruit development
Wine and table grape	1 l/ha	bunch elongation: post-fruit setting, grape swelling: from buckshot berry
Strawberry	100 ml/100 L of water	peduncle size and thickening: post-fruit setting
Fruit vegetables	100 ml/100 L of water	rachis/peduncle size and thickening: post-fruit setting
Leafy vegetables	100 ml/100 L of water	stay green of leaves throughout the entire crop cycle
Apple tree and watermelon	100 ml/100 L of water	Rachis/peduncle size and thickening: post-fruit setting
All crops	100-150 ml/100 L of water	recovery anti-stresses

# VigorGreen

## Plant metabolism promoter



VIGORGREEN is a cold-extracted concentrate of Canadian origin *Ascohyllum Nodosum* seaweed. This extraction allows to preserve the biologically active compounds (Polysaccharides, fucoidans, cytokines, auxins, gibberellins, betaines, alginates, mannitol, alginic acid etc) and ensure a more effective action on plants. The associated microelements (chelated iron DTPA, complexed boron and molybdenum) stimulate the metabolism of the plant by improving flowering, fruit set and uniformity of production. The present free amino acids with auxin action also stimulate fruit growth and production both qualitatively and quantitatively. Its use in the early stages (post-transplantation for horticultural plants, vegetative

recovery for fruit trees) improves the state of health of the plants and balances their growth.

### Benefits

- Promotes more balanced vegetative plant development
- Inverting action due to increased photosynthetic activity
- Improved Nutrient absorption;
- Increased blooming and greater fruit set
- Allows you to obtain fruits of greater size and better quality



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

Carbon (C) organic	6%
Iron (Fe) chelated with DTPA	0,3%
Boron (B) complexed with etalonamine,	0,5%
Molibdeno (Mo) soluble in water	0,04%
Mannitol	7 g/L
<b>ALSO ContAINS:</b>	
Free amino acids	5 g/kg
Polysaccharides	15%
Magnesium (MgO) soluble in water	2%
Alginic acid	15 g/kg
Fucoidans	15 g/kg
<b>PH</b>	<b>7,4</b>

### DENSITY

1,23 +/- 0,05 kg/dm<sup>3</sup>

Contains extracts of *Ascohyllum nodosum* algae.  
Color and density are indicative. For hazard warnings see page 154.

### DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Pome fruits	1-1,5 l/ha	From pre-flowering every 15 days
Stone fruits	1-2 l/ha	From pre-flowering every 15 days
Oil and table Olive	1-2 l/ha	From vegetative recovery onwards
Wine and table grapes	1-1,5 l/ha	From vegetative recovery onwards
Citrus fruits	1-1,5 l/ha	During the whole cycle
Actinidia (kiwi)	1-2 l/ha	From pre-flowering every 15 days
WalNut, hazel, chestNut	1-1,5 l/ha	From vegetative recovery onwards
Open field horticultural	1-2 l/ha	From post-transplant every 10-12 days
Horticultural in greenhouse	150-200 ml/100 liters	From post-transplant every 10-12 days
Leafy vegetables	1-1,5 l/ha	From post-transplant up to 10 days after harvest
Potato, carrot, onion, garlic, leek	1-1,5 l/ha	From post-emergency every 15 days



# Mycovix

## Resilient Roots



MYCOVIX has a high content of specific microorganisms that, thanks to their intense activity, are able to colonize the soil rapidly. This characteristic creates an ideal environment for root development (production of hormone-like substances, production of organic acids, and improvement of the soil's biological value) and an unfavorable environment for phytopathogens (space and Nutrient competition).

MYCOVIX is an innovative fertilizer that promotes the development of a healthy rhizosphere and stimulates the plant's natural defenses, improving its resistance, resilience, and vitality. MYCOVIX works synergistically to enhance the

growth and robustness of crops, helping maintain a healthy and vibrant soil ecosystem.

### Benefits

- Promotes the growth of beneficial microorganisms such as *Pochonia c.* and *Purpureocillium l.*;
- Increased root development;
- More fertile soil;
- Improved plant health;
- Enhanced Nutrient absorption.



**TYPE**  
Liquid



### PACKAGING



TRANSPARENT



### COMPOSITION

Mycorrhizae	0,1%
Rhizosphere bacteria (CFU/g)	2x10 <sup>7</sup>
Trichoderma	1x10 <sup>4</sup>
pH	7

### DENSITY

1,13 +/- 0,05 kg/dm<sup>3</sup>

The product does not contain genetically modified organisms and pathogenic organisms. Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	3-4 l/ha	1-2 spring applications 1-2 summer applications	fertigation
Open field horticultural	3-5 l/ha	From post-transplant (2-3 applications every 8-10 gg)	fertigation
Horticultural in greenhouse	400-500ml/1000m	From post-transplant every 8-12 gg	fertigation

## DIRECTION FOR USE

PREBIOTIC	DO NOT MIX WITH	NOTES
SuprEmo, Giove Bio Gold, Giove Bio, Verv N9, Naturblack	Stillages, phosphoric acid, nitric acid, products containing sulfuric acid, products containing copper, sulfur, bactericidal products, phosphites, and fungicides.	Space treatments with products containing phosphorus at least 7 days apart. Space any treatments with phosphites, fungicides, and copper-based products at least 10-15 days apart.

# A-Myco

## Balanced environment



A-MYCO is a fertilizer rich in beneficial microorganisms that stimulates plants' natural defenses, improving their resistance and vitality and creating a balanced environment ideal for healthy, vigorous growth.

A-MYCO is composed of a pool of specific microorganisms that act in a targeted and natural way, integrating into the agricultural ecosystem and sustainably supporting plant resilience.

### Benefits

- Promotes the development of beneficial microorganisms such as *Beauveria b.* and *Metharizium a.*;
- Healthy and lush foliage;
- Healthy and strong roots due to the metabolites produced by the microorganisms;
- Healthy and high-quality crops;
- It adapts well to nutritional strategies thanks to its anti-stress properties and resistance inducer.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Mycorrhizae</b>	<b>0,01%</b>
<b>Rhizosphere bacteria (CFU/g)</b>	<b>2x10<sup>7</sup></b>
<b>Trichoderma</b>	<b>1x10<sup>4</sup></b>
<b>pH</b>	<b>7</b>

### DENSITY

1,17 +/- 0,05 kg/dm<sup>3</sup>

The product does not contain genetically modified organisms and pathogenic organisms. Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	1-2,5 l/ha	Treat at the first signs of stress, repeat after 5-7 days	foliar
	2,5-3 l/ha		fertigation
Open field horticultural	1-2,5 l/ha	Treat at the first signs of stress, repeat after 5-7 days	foliar
	2,5-3,5 l/ha		fertigation
Horticultural in greenhouse	250-350 ml/1000m <sup>2</sup>	Treat at the first signs of stress, repeat after 5-7 days	foliar
	300-400 ml/1000m <sup>2</sup>		fertigation

## DIRECTION FOR USE

PREBIOTIC	DO NOT MIX WITH	NOTES
SuprEmo, Giove Bio Gold, Giove Bio, Verv N9, Naturblack	Stillages, phosphoric acid, nitric acid, products containing sulfuric acid, products containing copper, sulfur, bactericidal products, phosphites, and fungicides.	Space treatments with products containing phosphorus at least 7 days apart. Space any treatments with phosphites, fungicides, and copper-based products at least 10-15 days apart.



# Edafix

## Bacteria useful for soil and plants

EDAFIX contains rhizosphere bacteria capable of fixing nitrogen, promoting plant growth, and improving soil fertility. Mycorrhizal fungi increase the surface area explored by the roots of the plants they intercrop with, improving water and nutrient uptake.

*Azotobacter* spp. fixes atmospheric nitrogen into a form that plants can utilize, synthesizing growth-promoting substances and encouraging root development.

*Azospirillum* spp. forms a beneficial association with plant roots, fixing nitrogen and promoting the synthesis of substances such as cytokinins and auxins.



**TYPE**  
Water-soluble  
powder



### PACKAGING



### Benefits

- Improves nutrient uptake;
- Helps improve soil structure and the development of existing microbial flora;
- Increases overall plant growth;
- Improves crop productivity and quality.

### COMPOSITION

<b>Micorrize</b>	<b>&gt;10%</b>
<b>Rhizosphere bacteria (CFU/g)</b>	<b>5x10<sup>7</sup></b>
of which <i>Azospirillum</i> spp.	2,5x10 <sup>7</sup>
of which <i>Azotobacter</i> spp.	2,5x10 <sup>7</sup>
<b>pH</b>	<b>6</b>

### DENSITY

0,6 +/- 0,05 kg/dm<sup>3</sup>

The product does not contain genetically modified organisms and pathogenic organisms. Color and density are indicative


## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Cereals	0,40-0,50 kg/ha	Initial stages of vegetative development. Seed treatment	1-2 Foliar applications
Legumes	0,40-0,50 kg/ha	Initial stages of vegetative development	1-2 Foliar applications
Oleaginous	0,40-0,50 kg/ha	Initial stages of vegetative development. Seed treatment	1-2 Foliar applications
Industrial crops (beet, potato, sugar beet)	0,40-0,50 kg/ha	Initial stages of vegetative development	1-2 Foliar applications
Fruits trees	0,40-0,60 kg/ha	After fruit set. At the end of fruit swelling	2-3 Foliar applications 1-2 Fertigations

## DIRECTION FOR USE

PREBIOTIC	DO NOT MIX WITH	NOTES
SuprEmo, Giove Bio Gold, Giove Bio, Verv N9, Naturblack	Stillages, phosphoric acid, nitric acid, products containing sulfuric acid, products containing copper, sulfur, bactericidal products, phosphites, and fungicides.	Space treatments with products containing phosphorus at least 7 days apart. Space any treatments with phosphites, fungicides, and copper-based products at least 10-15 days apart.





## LEAF MACRO & MESOELEMENTS

Liquid fertilizers based on nitrogen, phosphorus, potassium and meso elements.

The LEAF line presents various formulations capable to satisfy the nutritional needs of crops in different phenological stages.

Each Leaf product contains high-quality raw materials and is formulated with adjuvants that improve and optimize absorption.

# Leaf N

## Gradual nitrogen nutrition

LEAF N contains gradual release nitrogen which guarantees constant and long-lasting nutrition over time, avoiding vegetative excesses. LEAF N also provides sulfur and boron which increase the protein content, the oil yield and the aromatic profile of the grapes.

### Benefits


- It increases production thanks to the presence of slow release nitrogen, which favors crop development and stay green;
- It helps to overcome the most difficult development phases in spring due to non-optimal conditions with an immediate greening effect;
- It improves fruit quality, increases the sugar level of cereals, increases the hectoliter weight of proteaginous plants and the protein content, thanks to the synergy between nitrogen and sulphur;
- It promotes flowering and fruit setting due to the presence of boron;
- Easy to use as it can be used both by foliar application and by fertigation.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>22%</b>
of which ureic nitrogen	11%
of which urea formaldehyde	7%
of which ammoniacal	4%
<b>Sulfur (SO<sub>2</sub>) soluble in water</b>	<b>11%</b>
<b>Boron (B) soluble in water</b>	<b>0,5%</b>
<b>pH</b>	<b>8</b>

Color and density are indicative

### DENSITY

1,25 +/- 0,05 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	2-3 l/ha	1-2 applications every 10-20 days from vegetative growth	foliar
Wine and table grape	2-3 l/ha	1-2 applications every 10-20 days from vegetative growth	foliar
Olive tree	2-3 l/ha	vegetative phase, budding, enlargement of the drupe	foliar
Straw cereals	3-4 l/ha	weeding at the end of tillering / beginning of raising, with flag leaf with fungicides	foliar
Extensive crops	3-4 l/ha	with weeding, fungicides and insecticides	foliar
Open field horticultural	2-3 l/ha	1-2 post-transplant applications	foliar
All crops	10-15 l/ha	at germination	fertigation

# Leaf P-Ca

## Specific formulation for size and quality

LEAF P-Ca contains a synergistic formulation of phosphorus and calcium bound together. LEAF P-Ca is effective at low dosages thanks to the high concentration of completely soluble phosphorus and calcium. LEAF P-Ca can be used both in the early stages, rooting and vegetative development, and in the stages of flowering, fruit set and development of the fruit (increasing size, uniformity, hardness and shelf life).

### Benefits

- Greater size and quality of the fruit (uniformity, hardness and shelf life);
- Synergistic formulation, bound and highly assimilable phosphorus and calcium;
- Low dosages thanks to the high concentration of phosphorus and calcium (P 23.6% - Ca 6%) completely soluble;
- Extremely versatile product that can be used in every phenological phase.



**TYPE**  
Liquid



### PACKAGING



TRANSPARENT



### COMPOSITION

<b>Nitrogen (N) nitric</b>	<b>3%</b>
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>23,6%</b>
<b>Calcium (CaO) soluble in water</b>	<b>6%</b>
<b>pH</b>	<b>0,5</b>

### DENSITY

1,37 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Stone fruits	4-5 l/ha	from fruit set:3 applications every 7-8 days	foliar
Pome fruits and kiwi	4-5 l/ha	from fruit set:3 applications every 10-14 days	foliar
Wine and table grape	5-6 l/ha	from fruit set:3 applications every 10-14 days	foliar
Olive tree	3 l/ha	from blossoming to fruit set	foliar
Open field horticultural	4-5 l/ha	beginning of fruit dev.:3-4 applications in 10-14 days	foliar
Horticultural in greenhouse	200-300 ml/hl	from fruit set:3-4 applications every 7-8 days	foliar
Potato	2 l/ha	10-15 days after emergence until tuberification	foliar
Alfa alfa	2,5 - 3 l/ha	in combination with herbicides and pesticides	foliar
All crops	10-15 l/ha	from the early stages of development	fertigation

# Leaf K

## Acid potassium for ripening

LEAF K stimulates ripening and uniforms it by increasing the color and the brix degree of the fruit. The particular formulation avoids overripening effects, maintaining the consistency of the fruit and the shelf-life. LEAF K can be used in association with pesticides as having a sub-acidic pH does not reduce their effectiveness and persistence.

### Benefits

- It stimulates ripening and makes it uniform increasing the color and the brix degree;
- It avoids over-ripening effects, maintaining fruit firmness and shelf-life;
- It increases yields by increasing the dry matter in the fruit;
- It can be used in association with pesticides as having a sub-acidic pH does not reduce its effectiveness and persistence.



**TYPE**  
Liquid



### PACKAGING



TRANSPARENT



### COMPOSITION

<b>Nitrogen (N) ureic</b>	<b>3%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>20%</b>
<b>pH</b>	<b>6,5</b>

### DENSITY

1,40 +/- 0,05 kg/dm<sup>3</sup>

### LOW CHLORINE

Color and density are indicative

## DOSAGES AND USES

CROPS	FOLIAR DOSAGES	PERIOD
Fruit trees	2-3 l/ha	2-3 applications da inizio veraison
Olive tree	2-3 l/ha	2-3 applications da fruit swelling in poi
Open field horticultural	2-3 l/ha	2-3 applications a partire da inizio veraison primi frutti
Horticultural in greenhouse	150-250 ml/hl	A partire da inizio veraison primi frutti
Wine and table grape	2-3 l/ha	2-3 applications da inizio veraison

# Leaf N-Fast

## Efficiency in nitrogen Nutrition

LEAF N-FAST is a highly efficient liquid nitrogen fertilizer. Nitrogen in the urea form complexed with lignosulfonates is rapidly absorbed and is 4 times more effective. LEAF N-FAST increases productivity and quality, stimulates photosynthetic activity, vegetative development and the rapid transformation of nitrogen into enzymes and proteins.

### Benefits



- Increase in productivity and quality, increase in photosynthetic activity, vegetative, development and rapid transformation of nitrogen into proteins;
- High efficiency nitrogen readily available;
- No risk of phytotoxicity even at high dosages;
- It improves the effectiveness of the combined products (fertilizers and/or pesticides), as thanks to the lignosulphonates, it has a wetting, tackifying and carrier effect.



**TYPE**  
Liquid



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) ureic</b>	<b>18%</b>
<b>Magnesium (MgO) water-soluble</b>	<b>4%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>8%</b>
<b>pH</b>	<b>7</b>

### DENSITY

1,25 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	5-10 l/ha	1-2 applications every 10-20 days from vegetative growth	foliar
Wine and table grape, hazelNut	5-10 l/ha	1-2 applications every 10-20 days from vegetative growth, in wine grapes apply pre-harvest to increase APA in musts	foliar
Olive tree	5-10 l/ha	Vegetative phase, cluster flowers, enlargement of the drupe	foliar
Straw cereals	10-20 l/ha	Weeding at the end of tillering / beginning of raising, with flag leaf and fungicides	foliar
Extensive crops	10-20 l/ha	With weeding, fungicides and insecticides	foliar
Open field horticultural	10-30 l/ha	1-2 applications during the vegetative development	foliar
Rice	3-5 l/ha	with Phytochemicals	foliar
All CROPS	15-30 l/ha	2-4 applications during all the cycle	FERTIGATION

# Leaf S-Quality

## Promotes quality and protein content

LEAF S-QUALITY contains nitrogen and sulfur totally available for the plant making it effective even at low dosages. LEAF S-QUALITY contains faster nitrogen in urea form and ammonia, more long-lasting over time, which guarantee a balanced development of the crop. LEAF S-QUALITY also provides sulfur which, when applied on the leaf, favors the accumulation of proteins, oil yield and the aroma of wine. In fertigation, it acidifies the rhizosphere and promotes the absorption of Nutrients

### Benefits

- It promotes quality by helping the accumulation of proteins, the yield in oil, the aroma of wine thanks to the balanced content of nitrogen and sulfur;
- It increases productivity by stimulating the plant to absorb the Nutrients present in the soil;
- It is efficient at low doses, since nitrogen and sulfur are totally available to the plant.



**TYPE**  
Liquid



### PACKAGING



TRANSPARENT



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>15%</b>
of which ammoniacal	10,4%
of which Nitrogen ureic	4,6%
<b>Sulfur (so<sub>3</sub>) soluble in water</b>	<b>57%</b>
<b>pH</b>	<b>8,5</b>

### DENSITY

1,33 +/- 0,05 kg/dm<sup>3</sup>

Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	5-10 l/ha	from vegetative growth	foliar
Table and Wine grape	3-4 l/ha	from post fruit set	foliar
Olive tree	4-5 l/ha	vegetative phase, inflorescence buds, enlargement of the drupe	foliar
Straw cereals	3-4 l/ha	weeding at the end of tillering / beginning of raising, at flag leaf/ear emergence with fungicides	foliar
Extensive crops	3-4 l/ha	with weeding, fungicides and insecticides	foliar
Open field horticultural	2-3 l/ha	1-2 post-transplant applications	foliar
All crops	15-20 l/ha	1-2 applications	fertigation

# HydroStar BTC

## Organo-mineral promoter of early stages of development

HYDROSTAR is a formulation composed of pure and highly soluble raw materials. HYDROSTAR contains a high percentage of organic nitrogen totally derived from plant extracts (enzymatic hydrolysis) with activating action for the plant and carrier for the mineral matrices present. The product has an acidic pH and is characterized by the absence of chlorides, sulphates, carbonates and urea nitrogen, which makes it usable in any vegetative phase and also for foliar.

These characteristics make HYDROSTAR a product that is effective at low dosages.

HYDROSTAR has an action of rooting and strengthening the plant while promoting flowering, thanks to the presence of pure extracts of the seaweed *Ascophyllum nodosum*.

### Benefits

- Better and faster planting;
- Better post-transplant recovery;
- Strengthening of the roots and collars of plants (increased resistance to environmental stress);
- Strengthening of the plant (closer flower stands);
- Improved flowering and fruit size.




### TYPE

Water-soluble powder



### PACKAGING



Contains extracts of *Ascophyllum nodosum* algae. Color and density are indicative

### COMPOSITION

<b>Nitrogen (N) total</b>	<b>13%</b>
of which Nitrogen organic	2%
of which Nitrogen (N) nitric	1,8%
of which Nitrogen ammoniacal	9,2%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>40%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>6%</b>
<b>Boron (B) soluble in water</b>	<b>0,03%</b>
<b>Iron (Fe) chelated with EDTA</b>	<b>0,04%</b>
<b>Manganese (Mn) chelated with EDTA</b>	<b>0,04%</b>
<b>Zinc (Zn) chelated with EDTA</b>	<b>0,04%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>
<b>pH</b>	<b>6</b>

### DENSITY

1,07 +/- 0,05 kg/dm<sup>3</sup>

### LOW CHLORINE

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR	PERIOD
Pome fruits	10-15 kg/ha	1,5-2 kg/ha	From vegetative growth until fruit set
Stone fruits	10-15 kg/ha	1,5-2,5 kg/ha	From vegetative growth until fruit set
Olive tree	10-15 kg/ha	2,5-3 kg/ha	From vegetative growth until fruit set
Wine and table grape	10-15 kg/ha	2-2,5 kg/ha	From vegetative growth until fruit set
Citrus trees	10-15 kg/ha	2-2,5 kg/ha	From vegetative growth until fruit set
Actinidia (kiwi)	10-20 kg/ha	2-2,5 kg/ha	From vegetative growth until fruit set
Nut, hazelnut, chestnut	10-15 kg/ha	1,5-2 kg/ha	From vegetative growth until fruit set
Open field horticultural	15-20 kg/ha	1-2 kg/ha	From post-transplant until first cluster of fruit set
Horticultural in greenhouse	1-2 kg/1000 metri	100-120 g/100 l of water	From post-transplant until third cluster of fruit set
OrLeafy vegetables	10-15 kg/ha	1,5-2 kg/ha	From post-transplant until half vegetative cycle
Potato, carrot, onion, garlic, leek	10-15 kg/ha	2-3 kg/ha	From post-emergence until tuber formation until tuber formation





# GRANULARS



## **NUECR4**

The efficient fertilizers

## **POWER**

Microgranular fertilizers

## **FERT**

Organo-mineral fertilizers

## **ACTIVE**

Mineral fertilizers

## **ORGANIC**

Organic fertilizers

## **SPECIALITY**

Slow release nitrogen fertilizers

# THE IMPORTANCE OF PHOSPHORUS FOR CEREAL FCP FERTILIZERS

CEREAL FCP places considerable importance on the type of phosphorus present in its fertilizers. The phosphate matrices used are treated to increase the percentage of water-soluble phosphorus.

Phosphorus is an essential element for plants, and its deficiency can cause serious consequences. It:

- promotes rooting, making the plant more resistant to disease and less susceptible to lodging;
- it concentrates in young tissue and is crucial for energy metabolism
- and in synthesis, breakdown, and transformation reactions;
- it is crucial for obtaining abundant and uniform flowering;
- it increases the speed of fruit ripening and improves their external quality.

**Phosphorus is characterized by poor mobility in the soil and low efficiency of use by plants** (on average, only 10-20% of the phosphorus added to the soil through fertilization is absorbed by the plant).

For this reason, also considering that phosphorus is retained by soil colloids, and therefore is not subject to losses through surface runoff, it is advisable, when possible, **to locate it close to the roots.**

- The solubility of phosphorus, and therefore its ease of absorption by plants, **depends on the soil pH:**

in highly acidic soils, phosphorus forms insoluble complexes with iron and aluminum hydroxides (phosphates Fe and Al);

in essentially neutral soils, it predominates in the form of monocalcium phosphate,  $\text{Ca}(\text{H}_2\text{PO}_4)_2$ , and dicalcium phosphate,  $\text{CaHPO}_4$ ;

in basic soils due to constitutional alkalinity (calcareous soils), the insoluble tricalcium phosphate,  $\text{Ca}_3(\text{PO}_4)_2$ , predominates;

in sodic soils, it predominates in the form of soluble sodium phosphate ( $\text{Na}_3\text{PO}_4$ ).

Phosphorus absorption is therefore favored in **essentially neutral soils**, while in acidic and basic soils it undergoes insolubilization (phosphorus retrogradation).

Thanks to the different types of phosphorus used, Cereal FCP products **are also particularly suitable for use on crops in alkaline soils**, however, rich in calcium carbonates and bicarbonates.



# THE EFFICIENCY OF FERTILIZERS

## NUE: NUTRIENT USE EFFICIENCY

Nutrient Use Efficiency (NUE) is a very important characteristic in plant nutrition, as plants absorb nutrients according to their characteristics and needs. NUE represents the amount of a fertilizer absorbed by the plant compared to the amount actually supplied through fertilization.

**The higher the NUE of a fertilizer, the better its environmental profile (fewer losses in the ecosystem) and the more efficient the agriculture (food production with less research and greater sustainability).**

NUE is influenced by: the operating environment (open field, greenhouse, soil-based or soil-less cultivation, etc.), the growing season, the type of crop (herbaceous or arboreal), genetics, and the quality of the inputs (fertilizers).

Through the adoption of different technologies, several CEREAL FCP products improve NUE, **particularly those from the NUECR4 line** and, to varying degrees, organo-mineral fertilizers or others like Mastercote.

## BIOACTIVATED FERTILIZERS

### COATING SYSTEM

CEREAL FCP is equipped with a modern coating plant to produce activated fertilizers **(one or more additives are added to the granules in post-granulation)**. This goal stems from CEREAL FCP's belief in the need to produce more efficient and environmentally friendly fertilizers.

Our coating process can be achieved using various additives:

- minerals, such as microelements
- organics, such as biostimulants
- innovative products, such as microorganisms

The use of high-quality raw materials, both liquid and powdered, allows us to enrich the granular product while preserving or increasing the performance of the original raw materials. This allows us to distribute special products to the soil, which would normally be applied through foliar application or fertigation.


The composition and concentration of the coating differ depending on the specific nutrient to be produced and its intended uses.

Activated fertilizers can be used on various extensive crops, fruit trees, and vegetables. Activating a granular fertilizer can offer several advantages, such as:

- **increasing nutrient use efficiency (NUE);**
- **reducing processing costs** by eliminating the need for multiple applications (use of inhibitors);
- introducing spores of microbial species that **solubilize nutrients locked** in the soil;
- **stimulating the root system** of plants with biostimulants (seaweed and amino acid cream).

The system is sized for maximum industrial flexibility and maximum precision in formulation.

### COATING : HOW TO BIOACTIVATE A GRANULAR FERTILIZER



# NUECR4

## THE EFFICIENT FERTILIZER



NUECR4 is the new family of granular which retain nutrients to release them gradually over time

The NUECR4 family is divided into two specific subcategories

- **NUECR4 PREMIUM:** class of products containing potassium sulphate for applications on crops sensitive to the presence of chlorine
- **NUECR4:** class of products for applications on crops more tolerant to the presence of chlorine

# LA TECNOLOGIA NUECR4




The NUECR4® technology in our granular fertilizers improves the NUE of nutrients (see “NUE: Nutrient Use Efficiency” on page 85).

## What is NUECR4® technology?

CR stands for “**Cation Retention**”: the fertilizer granules behave like clays in the soil, retaining cations and then releasing them over time.

## How does NUECR4® work?

Once distributed on the soil, the granules hydrate, and the nitrogen ( $\text{NH}_4^+$ ) and other cations (Mg, Ca, Mn, etc.) present are solubilized, binding, thanks to **NUECR4 Technology**, through cation retention. The release of the retained cations occurs gradually over time, allowing plants to utilize the elements more efficiently.



## Advantages of NUECR4®

- Increase in the Nutrient Use Efficiency (NUE)
- Slow release of elements
- Vigorous and constant growth of the plant since the earliest vegetative stages
- Increase in yield
- Greater sustainability

# Blurain



BLURAIN is a complex mineral NPK fertilizer with microelements obtained by granulation. BLURAIN is suitable for tree crops, small fruit, and open-field and greenhouse vegetables. The high soluble sulfur content improves nitrogen absorption. The product supports


plant nutrition beyond the early vegetative stages, promoting the development and renewal of the root system. Ideal for basal fertilization or band starter.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>10%</b>
of which Nitrogen Ammoniacal	8,5%
of which Nitrogen Ureic	1,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) solubile in citrato ammonico neutro e acqua</b>	<b>12%</b>
of which soluble in water	9%
<b>Potassium (K<sub>2</sub>O) solubile in water</b>	<b>16%</b>
<b>Magnesium (MgO)</b>	<b>2%</b>
of which soluble in water	1,5%
<b>Sulfur (SO<sub>3</sub>) solubile in water</b>	<b>20%</b>
<b>Boron (B)</b>	<b>0,01%</b>
<b>Iron (Fe)</b>	<b>1%</b>
<b>Zinc (Zn)</b>	<b>0,01%</b>

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,94 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Stone fruits	350-500 kg/ha	post-harvest/vegetative growth
Pome fruits	350-500 kg/ha	post-harvest/vegetative growth
Wine grape	350-500 kg/ha	post-harvest/vegetative growth
Table grape	400-600 kg/ha	post-harvest/vegetative growth
Kiwi	400-600 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	350-500 kg/ha	post-harvest/vegetative growth
Citrus trees	350-500 kg/ha	post-harvest/vegetative growth
Olive tree	350-600 kg/ha	post-harvest/vegetative growth
Estensive	200-300 kg/ha	localized at seeding
Horticultural in greenhouse	400-600 kg/ha	pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	400-600 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	400-600 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	300-500 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	350-550 kg/ha	pre-sowing, pre-transplant
Cabbages	350-550 kg/ha	pre-transplant
Garlic, onion, leek	350-550 kg/ha	pre-sowing, pre-transplant

# Control (slow release Nitrogen)

12.6.14+4CaO+2MgO+3SO<sub>3</sub>+0,02B+1Fe+0,01Zn




CONTROL is an NPK mineral fertilizer with highly soluble and available mineral matrices. It also contains 5% urea formaldehyde, which ensures greater nitrogen availability over time, ensuring a constant supply of the nutrient. CONTROL is characterized by a high soluble sulfur content, which improves nitrogen absorption.



**TYPE**  
Granular



## PACKAGING



LOW CHLORINE



The product supports plant nutrition beyond the early vegetative stages, promoting the development and renewal of the root system. CONTROL is a formulation suitable for horticultural crops, fruit trees, lawns, and golf courses.

## COMPOSITION

<b>Nitrogen (N) total</b>	<b>12%</b>
of which Nitrogen a lento rilascio (urea formaldeide)	5%
of which Nitrogen Ammoniacal	5,5%
of which Nitrogen Ureic	1,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) solubile in citrato ammonico neutro e acqua</b>	<b>6%</b>
of which soluble in water	5%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>14%</b>
<b>Calcium (CaO)</b>	<b>4%</b>
of which soluble in water	2%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>30%</b>
of which soluble in water	28%
<b>Boron (B)</b>	<b>0,02%</b>
<b>Iron (Fe)</b>	<b>1%</b>
<b>Zinc (Zn)</b>	<b>0,01%</b>


## DENSITY

0,90 kg/dm<sup>3</sup>

Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES SPAGLIO	DOSAGES LOCALIZZATO	PERIOD
Wine grape	350-450 kg/ha	175-225 kg/ha	post-harvest/vegetative growth
Table grape	350-550 kg/ha	175-275 kg/ha	post-harvest/vegetative growth
Kiwi	350-550 kg/ha	175-275 kg/ha	post-harvest/vegetative growth
Citrus trees	350-450 kg/ha	175-225 kg/ha	post-harvest/vegetative growth
Stone fruits	350-450 kg/ha	175-225 kg/ha	post-harvest/vegetative growth
Pome fruits	350-450 kg/ha	175-225 kg/ha	post-harvest/vegetative growth
Nursery and garden plants	350-550 kg/ha	175-275 kg/ha	pre-sowing, pre-transplant
Turf	30-50 gr/m <sup>2</sup>	15 - 25 kg/ha	pre-sowing/sowing
Nut, hazelnut, chestnut	350-450 kg/ha	175-225 kg/ha	post-harvest/vegetative growth
Open field horticultural and in greenhouse	300-500 kg/ha	150-250 kg/ha	pre-sowing, pre-transplant



# POWER

## STARTER EFFECT

POWER is the new family of micro-granular fertilizers, mineral and organo-mineral, obtained by granulation.

The micro-granule formulation allows their use at sowing directly in the furrow in contact with the seed, or at transplanting (starter technique). The small grain size of the Power fertilizers (0.5 - 1.7 mm for SuperPower, 1 - 2 mm for the remaining ones) and the low application dosages do not lead to phytotoxicity to the plant and provide nutrients starting from the early development stages.

The Power family offers a wide range of products that can be used both in conventional and organic farming, recommended for extensive and horticultural crops. The uniform granulometry and the hardness of the granule guarantee a high homogeneity of distribution through the micro-granulators present on the seed drills and transplanters.

The nitrogen present in Power fertilizers is of two types:

- ammoniacal: this is retained by the soil and it has low phytotoxicity for the plant,
- organic: vegetable or animal protein.

# SuperPower

**9.28+4CaO+15SO<sub>3</sub>**

SuperPower is a compound NP micro-granular mineral fertilizer (grain size 0.7 - 1.7 mm) with a high content of phosphorus, nitrogen, calcium and sulfur. The new formulation and the phosphorus / nitrogen ratio favours a pronounced starter effect by promoting the initial growth of the roots and allows it to be applied both in the seed furrow, in contact with the seed and in the transplant phase (starter technique).


This allows to reduce the dosages of use and a greater efficiency of use of the fertilizer. Particularly suitable for extensive and horticultural crop.



**TYPE**  
MicroGranular



**PACKAGING**



**GRAIN SIZE**  
95% between 0,5-1,7 mm

Color and density are indicative

**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>9%</b>
of which ammoniacal	9%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in neutral ammonium citrate and water</b>	<b>28%</b>
of which soluble in water	26%
<b>Calcium (CaO) soluble in water</b>	<b>4%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>15%</b>

**DENSITY**  
0,90 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Straw cereals	30-50 kg/ha	localized at seeding
Extensive crops	30-50 kg/ha	localized at seeding
Legumes	30-50 kg/ha	localized at seeding
Industrial crops (Tomato, potato, sugarbeet)	40-60 kg/ha	localized at seeding/transplant
Cotton	30-50 kg/ha	localized at seeding
Open field horticultural and in greenhouse	40-60 kg/ha	localized at seeding/transplant
Strawberry	30-50 kg/ha	localized at transplant

# SuperPower Humic

## 9.29+8CaO+18SO<sub>3</sub>+0,07B+0,1Zn+Acidi Umici



SuperPower Humic is a mineral micro-granular NP fertilizer with boron, zinc and humic acids (granulometry 0,7 - 1,7). Humic acids are from vegetal matrices that have undergone a long process of decay in soil, provide organic matter to the soil explored by the roots. It can be applied both in furrow, in contact with the seed, and during transplant (starter technique). This allows to reduce the


dosages thanks to an improved fertilizer use efficiency. It is recommended for field crops and horticultural.



**TYPE**  
MicroGranular



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>9%</b>
of which ammoniacal	9%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in neutral ammonium citrate and water</b>	<b>29%</b>
of which soluble in water	28%
<b>Calcium (CaO)</b>	<b>8%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>18%</b>
<b>Boron (B) soluble in water</b>	<b>0,07%</b>
<b>Zinc (Zn)</b>	<b>0,1%</b>

### GRAIN SIZE

95% between 0,5-1,7 mm

### DENSITY

0,90 kg/dm<sup>3</sup>

Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Extensive crops	30-50 kg/ha	localized at seeding
Industrial crops (Tomato, potato, sugarbeet)	40-60 kg/ha	localized at seeding/transplant
Strawberry	30-50 kg/ha	localized at transplant
Straw cereals	30-50 kg/ha	localized at seeding
Legumes	30-50 kg/ha	localized at seeding
Cotton	30-50 kg/ha	localized at seeding
Open field horticultural and greenhouse	40-60 kg/ha	localized at seeding/transplant

# SuperPower Plus

9.29+4CaO+15SO<sub>3</sub>+0,07B+0,1Zn



SuperPower Plus is a mineral micro-granular NP compound fertilizer with boron and zinc (granulometry 0,7 - 1,7 mm). The new formulation with boron and zinc promotes the plant growth, stimulates the root development, and provides several advantages:

- increases the nutrient use efficiency
- promotes root development and root density
- anticipates flowering and maturation

• is environmental friendly


It can be applied both in furrow, in contact with the seed, and during transplant (starter technique). This allows to reduce the dosages thanks to an improved fertilizer use efficiency. It is recommended for field crops and horticultural.



**TYPE**  
MicroGranular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>9%</b>
of which ammoniacal	9%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in neutral ammonium citrate and water</b>	<b>29%</b>
of which soluble in water	27%
<b>Calcium (CaO) soluble in water</b>	<b>4%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>15%</b>
<b>Boron (B) soluble in water</b>	<b>0,07%</b>
<b>Zinc (Zn)</b>	<b>0,1%</b>

**GRAIN SIZE**

95% between 0,5-1,7 mm

**DENSITY**

0,90 kg/dm<sup>3</sup>

Color and density are indicative

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Straw cereals	30-50 kg/ha	localized at seeding
Extensive crops	30-50 kg/ha	localized at seeding
Legumes	30-50 kg/ha	localized at seeding
Industrial crops (Tomato, potato, sugarbeet)	40-60 kg/ha	localized at seeding/transplant
Cotton	30-50 kg/ha	localized at seeding
Open field horticultural and in greenhouse	40-60 kg/ha	localized at seeding/transplant
Strawberry	30-50 kg/ha	localized at transplant

# SuperPower Extra

8.41+5CaO+6SO<sub>3</sub>+0.1B+0.02Cu+0.5Fe+0.5Zn



SuperPower Extra is a mineral micro-granular NP compound fertilizer. It has a high content of soluble phosphorus and can be absorbed in the first development stages of the plant. Boron, zinc and iron help to prevent chlorosis phenomena.

SuperPower Extra promotes root development, anticipates flowering and respects the environment. Its formulation


allows the application in furrow, in contract with the seed, or during transplant (starter technique), thus reducing dosages and improving the fertilizer efficiency.



**TYPE**  
MicroGranular



## PACKAGING



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>8%</b>
of which ammoniacal	8%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in neutral ammonium citrate and water</b>	<b>41%</b>
of which soluble in water	37%
<b>Calcium (CaO) soluble in water</b>	<b>5%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>6%</b>
<b>Boron (B) soluble in water</b>	<b>0,1%</b>
<b>Copper (Cu) soluble in water</b>	<b>0,02%</b>
<b>Iron (Fe)</b>	<b>0,5%</b>
<b>Zinc (Zn)</b>	<b>0,5%</b>

## GRAIN SIZE

95% between 0,5-1,7 mm

## DENSITY

0,90 kg/dm<sup>3</sup>

Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Industrial crops (Tomato, potato, sugarbeet)	35-50 kg/ha	localized at seeding/at transplant
Strawberry	30-50 kg/ha	localized at transplant
Open field horticultural and in greenhouse	35-50 kg/ha	localized at seeding/at transplant
Extensive crops	30-50 kg/ha	localized at seeding
Straw cereals	30-50 kg/ha	localized at seeding
Legumes	30-50 kg/ha	localized at seeding
Cotton	30-50 kg/ha	localized at seeding

# Power BioAger

## 4.6.10+10CaO+2MgO+16SO<sub>3</sub>+3Fe+12C-Org



Power BioAger is a compound micro-granular organo-mineral NPK fertilizer obtained by granulation from high quality raw materials allowed in organic farming. Its composition makes Power BioAger ideal for fruit trees, Wine grape, olive trees, and horticultural crops. The reduced granulometry allows distributing Power BioAger both with spreaders, microgranulators, and transplanting machines (starter technique). The C/N ratio equals to 3 allows to


microorganisms into the soil to transform the organic nitrogen in nitrogne that can be aborbed by the plant, in order to avoid the nitrogen to kept by the soil-plant system.



**TYPE**  
MicroGranular



**PACKAGING**



**COMPOSITION**

Nitrogen (N) organic	4%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) soluble in mineral acids of which at least 55% soluble in formic acid	2%
Potassium (K <sub>2</sub> O) soluble in water	10%
Calcium (CaO)	10%
Magnesium (MgO) soluble in water	2%
Sulfur (SO <sub>3</sub> ) soluble in water	16%
Iron (Fe)	3%
Organic carbon	12%

**GRAIN SIZE**  
95% between 1-2 mm

**DENSITY**  
0,88 kg/dm<sup>3</sup>

Color and density are indicative

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Open field horticultural and in greenhouse	50-80 kg/ha	Localized at seeding/at transplant
Industrial crops (Tomato, potato, )	50-80 kg/ha	Localized at seeding/at transplant
Strawberry	50-80 kg/ha	Localized at transplant
Cotton	40-70 kg/ha	Localized at seeding
Straw cereals	40-70 kg/ha	Localized at seeding
Extensive crops	40-70 kg/ha	Localized at seeding
Legumes	40-70 kg/ha	Localized at seeding

# Power BioMaster

## 6.5.13+8CaO+16SO<sub>3</sub>+3Fe+20C-org

bioagricert 



Power BioMaster is an NPK organo-mineral compound fertilizer micro-granular obtained by granulation from noble raw materials allowed in organic farming. The ratio of nitrogen, phosphorus, potassium and sulfur makes it ideal for fertilizing fruit crops, Wine grape and vegetables. The


conformation of the granule and its granulometry make it suitable for distribution both with fertilizer spreaders and with distributors present in seeders and transplanters. Ideal for applying the starter technique at transplant or at sowing.



**TYPE**  
MicroGranular



### PACKAGING



### COMPOSITION

Nitrogen (N) organic	6%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) soluble in mineral acids of which at least 55% soluble in formic acid 2%	5%
Potassium (K <sub>2</sub> O) soluble in water	13%
Sulfur (SO <sub>3</sub> ) soluble in water	16%
Calcium (C <sub>a</sub> O)	8%
Iron (Fe)	3%
Organic carbon	20%

**GRAIN SIZE**  
95% between 1-2 mm

**DENSITY**  
0,84 kg/dm<sup>3</sup>

Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Industrial crops (Tomato, potato, sugarbeet)	50-80 kg/ha	localized at seeding/at transplant
Open field horticultural and in greenhouse	50-80 kg/ha	localized at seeding/at transplant
Strawberry	50-80 kg/ha	localized at transplant
Cotton	40-70 kg/ha	localized at seeding
Straw cereals (grano, orzo, segale)	40-70 kg/ha	localized at seeding
Extensive crops	40-70 kg/ha	localized at seeding
Legumes	40-70 kg/ha	localized at seeding

# Power BioNascor

## 5.10+12CaO+8SO<sub>3</sub>+6Fe+12C-Org

bioagricert 



Power BioNascor is an NP micro-granular organic fertilizer obtained by granulation from a mixture of high quality organic matrices allowed in organic farming. The organic substance inside is rich in amino acids, humic and fulvic acids that help the absorption of nutrients and


restore the biological fertility of the soil. Its granulometry makes it suitable for the starter technique at transplant or sowing. Power BioNascor is ideal for fertilizing cereals, vegetables and orchards.



**TYPE**  
MicroGranular



### PACKAGING



### COMPOSITION

Nitrogen (N) organic	5%
Phosphorus (P <sub>2</sub> O <sub>5</sub> )	10%
Potassium (K <sub>2</sub> O) soluble in water	1,5%
Calcium (C <sub>a</sub> O)	12%
Sulfur (SO <sub>3</sub> ) soluble in water	8%
Iron (Fe)	6%
Organic carbon	12%

### GRAIN SIZE

95% between 1-2 mm

### DENSITY

0,84 kg/dm<sup>3</sup>

Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	40-70 kg/ha	localized at seeding
Extensive crops	40-70 kg/ha	localized at seeding
Legumes	40-70 kg/ha	localized at seeding
Industrial crops (Tomato, potato, sugarbeet)	50-80 kg/ha	localized at seeding/at transplant
Cotton	40-70 kg/ha	localized at seeding
Open field horticultural and in greenhouse	50-80 kg/ha	localized at seeding/at transplant
Strawberry	50-80 kg/ha	localized at transplant



# FERT

## THE BENEFITS OF THE TWO WORLDS



Family of NPK and NP organo-mineral granular fertilizers that are characterized by the presence of slow release protein nitrogen and humic acids.

- Fert granular fertilizers are obtained by granulation, therefore each granule contains within it all the nutritional elements present in the formula. This guarantees a high uniformity of distribution of the nutrients in the soil, improving their effectiveness for the plants.
- The raw materials used are high quality organic matrices, containing Humic Acids which:
  - bring organic matter to the soil
  - promote roots' nutrient absorption
  - protect against retrogradation (phosphorus insolubility).
- 
- The organic nitrogen contained in Fert fertilizers guarantees a gradual release by microorganisms present in soil, facilitating the radical absorption. Furthermore, Fert fertilizers contain soluble and assimilable phosphorus, assisted by a physiologically acid reaction. All formulations are created with a high soluble sulfur content. Their homogeneous granulometry guarantees good flexibility of use both in the open field and during seeding.
- 
- Fert family is divided into two sub-categories:
  - **FERT PREMIUM:** class of products containing potassium sulphate, for application on crops sensitive to the presence of chlorine
  - **FERT:** class of products for application on crops that are more tolerant to the presence of chlorine.
-

# ORGANIC MATTER, HUMUS AND HUMIC ACIDS IN OUR FERTILIZERS


The soil is formed by a solid component (sand, silt, clay and organic matter), a liquid component (water) and a gaseous component (oxygen and carbon dioxide). Among the solid compounds, the organic matter plays an important role since it regulates the fundamental properties of the soil (see figure).

The organic matter is formed by a mix of compounds among which the most relevant are:

- **the living biomass** (plant roots, microorganisms, insects, etc)
- **the dead biomass** (plant and animal wastes in decay)
- **humus** (resulting product of the activity of microorganisms present in the soil)

Humus is the most noble and important constituent in the organic matter. It is the most stable (and more difficult to degrade) and reactive component and, for this reason, all fundamental properties of the soil depend on it. Humus in turn contains humic and fulvic acids. The first ones are stable insoluble compounds, while the second ones are soluble products. This characteristic allows to extract and separate the two fundamental components of the organic matter and to obtain solid or liquid concentrates.

**The fertilizers of the FERT line contain humic acids from a vegetal matrix which has undergone a long process of decay in the soil.**



The FERT line fertilizers contain humic acids from plant matter that has undergone a long decomposition process in the soil.

# Granoro

**10.20+4CaO+20SO<sub>3</sub>+0,01B+0,03Zn+7,5C-org 2N org**



Granoro is an organo-mineral NP compound fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value and a soluble phosphorus content. The presence of humic acids allows a better assimilation of nutrients. Its high content

of soluble sulphur improves the protein content in cereal crops.


It is ideal for pre-sowing applications on winter cereals



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>10%</b>
of which organic	2%
of which ammoniacal	8%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>20%</b>
of which soluble in neutral ammonium citrate and water	16%
of which soluble in water	14%
<b>Calcium (CaO) soluble in water</b>	<b>4%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>20%</b>
<b>Zinc (Zn)</b>	<b>0,03%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,90 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	200-400 kg/ha	pre-sowing/seeding
Extensive	200-400 kg/ha	pre-sowing/seeding
Legumes	200-400 kg/ha	pre-sowing/seeding
Industrial crops (Tomato, Potato, sugarbeet)	300-500 kg/ha	pre-sowing, pre-transplant, anche localizzato
alfa alfa / fodder crops	300-500 kg/ha	pre-sowing/seeding
Open field horticultural and in greenhouse	200-300 kg/ha	pre-sowing, pre-transplant

# Flex

**6.10.18+3CaO+2MgO+12SO<sub>3</sub>+0,01B+0,03Zn+7,5C-org 2N org**



Flex is an organo-mineral NPK compound fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value. The balanced ratio among nutrients is ideal for the fertilization of alkaline soils. 2% of slow-release organic nitrogen

guarantees a constant absorption over time. Phosphorus is largely soluble, therefore it is assimilated by the plant more easily.


It is recommended for pre-planting applications on extensive and oil crops.



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>6%</b>
of which organic	2%
of which ammoniacal	4%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>10%</b>
of which soluble in neutral ammonium citrate and water	8%
of which soluble in water	4%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>18%</b>
<b>Calcium (C<sub>3</sub>O) soluble in water</b>	<b>3%</b>
<b>Magnesium (MgO)</b>	<b>2%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>12%</b>
<b>Zinc (Zn)</b>	<b>0,03%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,93 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	600-800 kg/ha	post-harvest/vegetative growth
Wine grapeE	400-600 kg/ha	post-harvest/vegetative growth
Table grape	600-800 kg/ha	post-harvest/vegetative growth
Citrus trees	600-800 kg/ha	post-harvest/vegetative growth
Straw cereals	200-400 kg/ha	pre-sowing/seeding
Legumes	200-400 kg/ha	pre-sowing/seeding
Industrial crops (Tomato, potato, sugarbeet)	600-800 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	600-800 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	600-800 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	500-700 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	600-800 kg/ha	pre-sowing, pre-transplant
Cabbages	400-600 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	600-800 kg/ha	Pre-sowing, pre-transplant

# Ortofrutto Special

12.6.6+3CaO+26SO<sub>3</sub>+0,01B+7,5C-Org 1N-Org



Ortofrutto Special is an organo-mineral NPK compound fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value. 1% of slow-release organic nitrogen guarantees a constant absorption over time, and phosphorus is soluble, therefore it is assimilated by the plant more easily. Boron favours the

pollination and fertility of flowers, while the presence of humic acids allows a more rapid and efficient translocation of nutrients in the plant and provide organic matter to the soil.


It is recommended to improve the quality and productivity of fruit trees and horticultural crops.



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>12%</b>
of which organic	1%
of which ammoniacal	11%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>6%</b>
of which soluble in neutral ammonium citrate and water of which soluble in water	5%
	4%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>6%</b>
<b>Calcium (CaO) soluble in water</b>	<b>3%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>26%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative.  
For hazard warnings see page 154.

## DENSITY

0,84 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	450-650 kg/ha	post-harvest/vegetative growth
Citrus trees	450-650 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	450-650 kg/ha	post-harvest/vegetative growth
Olive oil	450-650 kg/ha	post-harvest/vegetative growth
Table olive	450-650 kg/ha	post-harvest/vegetative growth
Horticultural in greenhouse	400-600 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	350-550 kg/ha	pre-sowing, pre-transplant
Cabbages	350-550 kg/ha	Pre-sowing, pre-transplant

# Ortofrutto Special Top

14.5.5+2CaO+30SO<sub>3</sub>+0,01B+7,5C-Org 1N-Org



Ortofrutto Special Top is an organo-mineral NPK compound fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value. It has a high content of soluble phosphorus, easy to be absorbed by the plant. The presence of humic acids guarantees a better assimilation of nutrients and provides

organic matter to the soil. Boron favours pollination and the fertility of flowers.


It is ideal to improve the quality and productivity of fruit trees and horticultural crops.



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>14%</b>
of which organic	1%
of which ammoniacal	13%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>5%</b>
of which soluble in neutral ammonium citrate and water of which soluble in water	4%
	3%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>5%</b>
<b>Calcium (CaO) soluble in water</b>	<b>2%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>30%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,86 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	450-650 kg/ha	post-harvest/vegetative growth
Citrus trees	450-650 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	450-650 kg/ha	post-harvest/vegetative growth
Olive oil	450-650 kg/ha	post-harvest/vegetative growth
Table olive	450-650 kg/ha	post-harvest/vegetative growth
Horticultural in greenhouse	400-600 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	350-550 kg/ha	pre-sowing, pre-transplant
Cabbages	350-550 kg/ha	pre-sowing, pre-transplant



# Master Plus

**10.8.15+3CaO+2Mgo+26SO<sub>3</sub>+0,01B+0,03Zn+7,5C-org 2N org**



Master Plus is an organo-mineral NPK compound fertilizer obtained by granulation with highly soluble mineral matrices. It has a slightly acidic pH that facilitates the mobility of nutrients into the soil. The synergy between nitrogen and sulphur increases the protein content in crops.


It is ideal for pre-sowing applications on cereal and oil crops, fodder plants, and fruit trees.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>10%</b>
of which organic	2%
of which ammoniacal	8%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>8%</b>
of which soluble in neutral ammonium citrate and water	7%
of which soluble in water	6%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>15%</b>
<b>Calcium (CaO) soluble in water</b>	<b>3%</b>
<b>Magnesium (MgO)</b>	<b>2%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>26%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Zinc (Zn)</b>	<b>0,03%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,90 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits (Apple tree, Pear tree)	300-500 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	500-700 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	400-600 kg/ha	post-harvest/vegetative growth
Citrus trees	600-800 kg/ha	post-harvest/vegetative growth
Small fruits	300-500 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	400-600 kg/ha	post-harvest/vegetative growth
Olive oil	400-600 kg/ha	post-harvest/vegetative growth
Table olive	400-600 kg/ha	post-harvest/vegetative growth
Industrial crops (Tomato, Potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	400-600 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	500-700 kg/ha	pre-sowing, pre-transplant

# Olive tree

**12.6.6+3CaO+26SO<sub>3</sub>+0.01B+7,5C-Org 1N-Org**



Olivo is an organo-mineral NPK compound fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value. 1% of slow-release organic nitrogen guarantees a constant absorption over time. It has a high content of soluble phosphorus, easy to be absorbed by the plant. Boron favours pollination and


the fertility of flowers. The presence of humic acids allows a more rapid and efficient translocation of nutrients in the plant and provide organic matter to the soil. It is ideal to improve the quality and productivity of fruit trees and olive trees.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>12%</b>
of which organic	1%
of which ammoniacal	11%
<b>Phosphorus (P<sub>2</sub>O<sub>3</sub>) total</b>	<b>6%</b>
of which soluble in neutral ammonium citrate and water	5%
of which soluble in water	4%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>6%</b>
<b>Calcium (CaO) soluble in water</b>	<b>3%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>26%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,84 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Olive oil	450-650 kg/ha	post-harvest/vegetative growth
Table olive	450-650 kg/ha	post-harvest/vegetative growth
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	450-650 kg/ha	post-harvest/vegetative growth
Citrus trees	450-650 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	450-650 kg/ha	post-harvest/vegetative growth

# Dual Band

**7.22+3CaO+16SO<sub>3</sub>+0,01Zn+7,5C-Org 1,5N-Org**



Dual Band is a NP organo-mineral fertilizer activated by humic and fulvic acids and obtained through granulation of selected raw materials.

Dual Band contains organic nitrogen of amino-acidic origin, zinc, calcium and sulphur, which all favour an optimal sprouting of the seed and development of the

root system. The presence of humic and fulvic acids have a biostimulant action and, moreover, guarantees nutrient availability for the crop limiting processes of retrogradation of phosphorus and nitrogen leaching.


Dual Band can be applied to all crops, in particular during the stage of pre-sowing of winter cereal crops.



**TYPE**  
Granular



## PACKAGING



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>7%</b>
of which organic	1,5%
of which ammoniacal	5,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>22%</b>
of which soluble in neutral ammonium citrate and water	15%
of which soluble in water	9%
<b>Calcium (C<sub>a</sub>O) soluble in water</b>	<b>3%</b>
<b>Sulfur (SO<sub>2</sub>) soluble in water</b>	<b>16%</b>
<b>Zinc (Zn)</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative.  
For hazard warnings see page 154.

**DENSITY**  
0,95 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	250-450 kg/ha	pre-sowing/seeding
Extensive crops	250-450 kg/ha	pre-sowing/seeding
Legumes	250-450 kg/ha	pre-sowing/seeding
Industrial crops (Tomato, potato, sugarbeet)	350-550 kg/ha	pre-sowing, pre-transplant, anche localizzato
Alfa alfa / fodder crops	350-550 kg/ha	pre-sowing/seeding
Open field horticultural and in greenhouse	250-350 kg/ha	pre-sowing, pre-transplant

# TrioStart

**8.15.8+4CaO+16SO<sub>3</sub>+0,01B+0,003Zn+7,5C-Org 2N-Org**



TrioStart is an organo-mineral NPK complex fertilizer rich in phosphorus which is totally assimilable.

The organic matrix which derives from hydrolyzed gelatin (rich in amino acids) improves the structure of the soil and ensures a slow-release nitrogen nutrition. The phosphorus present is not retrogradable because it is bound to a


humic matrix (humic acids) which allows its complete assimilation. It is particularly suitable for the stimulation of the root system, to strengthen the plant and, in the meantime, to reintegrate soil fertility.



**TYPE**  
Granular



**PACKAGING**



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>8%</b>
of which organic	2%
of which ammoniacal	6%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>15%</b>
of which soluble in neutral ammonium citrate and water	13%
of which soluble in water	11%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>8%</b>
<b>Calcium (CaO) soluble in water</b>	<b>4%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>16%</b>
<b>Boron (B)</b>	<b>0,01%</b>
<b>Zinc (Zn)</b>	<b>0,03%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,85 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Actinia (kiwi)	600-800 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	600-800 kg/ha	post-harvest/vegetative growth
Industrial crops (Tomato, Potato, sugarbeet)	600-800 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages	300-500 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	300-500 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	400-600 kg/ha	pre-sowing, pre-transplant

# Vinfrutto

**8.6.14+3CaO+2MgO+30SO<sub>3</sub>+0,01B+0,5Fe+0,01Zn+7,5C-org 2N org**



Vinfrutto is an NPK complex organo-mineral fertilizer containing microelements. It is obtained by granulation and it has a low chlorine content. Its formulation allows a complete and balanced intake of essential nutrients for plants. It contains:

- Highly soluble phosphorus (protected by humic acids), therefore more easily assimilable
- Soluble sulphur, calcium, magnesium, boron, iron and zinc which guarantee a complete nutrition.
- Vinfrutto is mostly recommended for autumn and/or spring fertilization of fruit trees.


- 2% organic nitrogen: slow-release nitrogen which guarantees an absorption over time (excellent carbon/nitrogen ratio which stimulates microbial activity in the soil)



**TYPE**  
Granular



**PACKAGING**



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>8%</b>
of which organic	2%
of which ammoniacal	6%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>6%</b>
of which soluble in neutral ammonium citrate and water	5%
of which soluble in water	3%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>14%</b>
<b>Calcium (CaO) soluble in water</b>	<b>3%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>30%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Iron (Fe)</b>	<b>0,5%</b>
<b>Zinc (Zn) soluble in water</b>	<b>0,01%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

### DENSITY

0,99 kg/dm<sup>3</sup>

**LOW CHLORINE**

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Actinia (kiwi)	600-800 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	600-800 kg/ha	post-harvest/vegetative growth
Citrus trees	600-800 kg/ha	vegetative growth
Small fruits	300-500 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	600-800 kg/ha	post-harvest/vegetative growth
Olive oil	400-600 kg/ha	vegetative growth
Table olive	400-600 kg/ha	vegetative growth
Horticultural in greenhouse	600-800 kg/ha	pre-sowing, pre-transplant
Industrial crops	600-800 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	500-700 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	500-700 kg/ha	pre-sowing, pre-transplant
Cabbages	500-700 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	500-700 kg/ha	pre-sowing, pre-transplant

# Ortoplus

**8.16.8+5CaO+25SO<sub>3</sub>+0,01B+0,02Zn+7,5C-Org 2N-Org**



Ortoplus is an organo-mineral NPK compound fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value. It has a high content of soluble phosphorus, therefore easily absorbed by the plant. Moreover, the presence of humic acids allows a better assimilation of nutrients.


It contains 2% of slow-release organic nitrogen which guarantees a constant absorption over time



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>8%</b>
of which organic	2%
of which ammoniacal	6%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>16%</b>
of which soluble in neutral ammonium citrate and water of which soluble in water	14%
	12%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>8%</b>
<b>Calcium (CaO) soluble in water</b>	<b>5%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>25%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Zinc (Zn) soluble in water</b>	<b>0,02%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
1,1 kg/dm<sup>3</sup>

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	600-800 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	600-800 kg/ha	post-harvest/vegetative growth
Industrial crops (Tomato, potato, sugarbeet)	600-800 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages	300-500 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	300-500 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	400-600 kg/ha	pre-sowing, pre-transplant

# Jolly

**6.10.20+3CaO+22SO<sub>3</sub>+7,5C-Org 2N-Org**



Jolly is an organo-mineral NPK fertilizer obtained by granulation. It contains both organic and mineral matrices with a high nutritional value. Its high sulphur content improves the quality of the organoleptic aspects of crops. It has a high potassium sulphate content. It is ideal for fall


and spring fertilization of fruit trees. It contains 2% of slow-release organic nitrogen which guarantees a constant absorption over time. Phosphorus is soluble, therefore easily assimilable.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>6%</b>
of which organic	2%
of which ammoniacal	4%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>10%</b>
of which soluble in neutral ammonium citrate and water of which soluble in water	8%
	6%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>20%</b>
<b>Calcium (CaO) soluble in water</b>	<b>3%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>22%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
1 kg/dm<sup>3</sup>

**LOW CHLORINE**

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	400-600 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	600-800 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	600-800 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	400-600 kg/ha	post-harvest/vegetative growth
Citrus trees	600-800 kg/ha	post-harvest/vegetative growth
Industrial crops (Tomato, potato, sugarbeet)	600-800 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	600-800 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	600-800 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	500-700 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	600-800 kg/ha	pre-sowing, pre-transplant
Cabbages	400-600 kg/ha	pre-sowing, pre-transplant
Tobacco	600-800 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	600-800 kg/ha	pre-sowing, pre-transplant

# Vinfrutto Star

10.10.15+2MgO+28SO<sub>3</sub>+7,5C-Org 2N-Org



Vinfrutto Star is an NPK organo-mineral complexed fertilizer, which is balanced and complete. The high title and the low chlorine content make the product of elevated quality. The ratio among nutrients ensures to the plant a complete and constant nutrition during the whole crop cycle. Vinfrutto Star is ideal for every crop, especially for the fertilization of post-harvest/autumn – winter fruit crops.

Vinfrutto star contains:


- 2% of organic nitrogen of high quality, rich in amino acids and polypeptides which allow a gradual absorption over time;
- Assimilable phosphorus and non retrogradable in the soil;
- Magnesium and sulphur with a greening and acidifying action;
- Humic acids which improve the structure of the soil and promote the absorption of nutrients.



**TYPE**  
Granular



**PACKAGING**



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>10%</b>
of which organic	2%
of which ammoniacal	8%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>10%</b>
of which soluble in neutral ammonium citrate and water	9%
of which soluble in water	8%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>15%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>28%</b>
<b>Carbon (C) organic</b>	<b>7,5%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,97 kg/dm<sup>3</sup>

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits (pesco, nettarina, percoco, albicocco, susino, mandorlo, ciliegio)	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits (Apple tree, Pear tree)	400-600 kg/ha	post-harvest/vegetative growth
Wine grape	350-600 kg/ha	post-harvest/vegetative growth
Table grape	500-700 kg/ha	post-harvest/vegetative growth
Kiwi	500-700 kg/ha	post-harvest/vegetative growth
Horticultural in greenhouse	500-700 kg/ha	pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	600-800 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	400-600 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	500-700 kg/ha	pre-sowing, pre-transplant
Cabbages	400-600 kg/ha	pre-transplant
Garlic, onion, leek	400-600 kg/ha	pre-sowing, pre-transplant





# ACTIVE LOVE FOR YOUR CROP

Family of P-NP-NPK granular mineral fertilizers with low chlorine content which are characterized by high solubility and availability of nutrients.

The Active family fertilizers contain phosphorus and other highly soluble nutrients. Their homogeneous granulometry guarantees a high uniformity of distribution.

They guarantee, thanks to the combination of nitrogen and sulfur, a greater protein content especially in cereal crops and a better absorption of nitrogen in all the others, thus limiting the losses (especially those for leaching).

All Active fertilizers have a physiologically acidic pH that increase the availability of phosphorus and microelements, which are more easily absorbed by the roots.

Active family is divided into ACTIVE PREMIUM and ACTIVE:

- **ACTIVE PREMIUM** line with sulphate potassium for applications in chlorine-sensitive crops;
- **ACTIVE** line for applications on more tollerant crops.

# Cereaphos

**0.40.0+12CaO+5SO<sub>3</sub>+0,01B+0,01Zn**



Cereaphos is a mineral fertilizer obtained by granulation with a high solubility and availability phosphatic matrix. The presence of sulphur increases the organoleptic aspects of crops, while boron favours flowering and fruit set by ensuring a more stable production. Zinc is important as it is a forerunner of many physiological reactions of plants and


it is also a cofactor in phosphorus absorption. Cereaphos contains humic acids. It is indicated for localized fertilization at sowing or transplant.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>40%</b>
of which soluble in water	36%
of which soluble in neutral ammonium citrate and water	40%
<b>Calcium (CaO) soluble in water</b>	<b>12%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>5%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Zinc (Zn) soluble in water</b>	<b>0,01%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,99 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Industrial crops (Tomato, potato, sugarbeet)	300-500 kg/ha	post-harvest/vegetative growth
Extensive crops	300-400 kg/ha	post-harvest/vegetative growth
Open field horticultural and in greenhouse	300-400 kg/ha	post-harvest/vegetative growth
Straw cereals	200-400 kg/ha	post-harvest/vegetative growth
Legumes	200-400 kg/ha	post-harvest/vegetative growth

# Universal Up

**9.30+6CaO+14SO<sub>3</sub>+0,01B+0,03Zn**



Universal Up is an NP mineral compound fertilizer activated with humic acids. It contains different forms of nitrogen (ammoniacal, ureic) and presents a high content of soluble phosphorus. Humic acids facilitate the absorption of nutrients and protect phosphorus from retrogradation

(insolubility).


It is indicated for fruit trees and extensive crops.



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>9%</b>
of which Nitrogen ammoniacal	7%
of which Nitrogen ureic	2%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>30%</b>
of which soluble in water	26%
of which soluble in neutral ammonium citrate and water	28%
<b>Calcium (CaO) soluble in water</b>	<b>6%</b>
<b>Sulfur (SO<sub>2</sub>) soluble in water</b>	<b>14%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Zinc (Zn)</b>	<b>0,03%</b>

Contains humic acids. Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,90 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	200-400 kg/ha	pre-sowing/seeding
Extensive crops	300-400 kg/ha	pre-sowing/seeding
Legumes	200-400 kg/ha	pre-sowing/seeding
Industrial crops (Tomato, potato, sugarbeet)	300-500 kg/ha	pre-sowing, pre-transplant, also localized
Open field horticultural and in greenhouse	300-400 kg/ha	pre-sowing, pre-transplant



# Land Plus

**10.8.15+3CaO+2MgO+3ISO<sub>3</sub>+0,01B+0,01Cu+0,001Mo+0,01Zn**



Land Plus is an NPK mineral compound fertilizer with activator, obtained by granulation. It contains high-solubility and availability mineral matrices. The high amount of nitrogen in different forms (ammoniacal and ureic) and the synergy with sulphur improve the protein content in several crops. Its balanced formulation and

the presence of calcium, magnesium, sulphur and microelements ensure a complete nutrition to the plant. It contains humic acids.


Land Plus is indicated for pre-sowing applications on cereal crops, oil crops and fruit trees.



**TYPE**  
Granular



## PACKAGING



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>10%</b>
of which Nitrogen Ammoniacal	8%
of which Nitrogen Ureic	2%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>8%</b>
of which soluble in water	5%
of which soluble in neutral ammonium citrate and water	6%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>15%</b>
<b>Calcium (CaO) soluble in water</b>	<b>3%</b>
<b>Magnesium (MgO)</b>	<b>2%</b>
of which soluble in water	1,5%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>31%</b>
of which soluble in water	28%
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Copper (Cu)</b>	<b>0,01%</b>
<b>Molibdeno (Mo)</b>	<b>0,001%</b>
<b>Zinc (Zn)</b>	<b>0,01%</b>

Color and density are indicative. For hazard warnings see page 154.

## DENSITY

0,90 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	400-600 kg/ha	post-harvest/vegetative growth
Kiwi	400-600 kg/ha	post-harvest/vegetative growth
Nut, hazelnut, chestnut	400-600 kg/ha	post-harvest/vegetative growth
Citrus trees	400-600 kg/ha	post-harvest/vegetative growth
Stone fruits	300-500 kg/ha	post-harvest/vegetative growth
Pome fruits	300-500 kg/ha	post-harvest/vegetative growth
Olive oil	300-500 kg/ha	vegetative growth
Table olive	300-500 kg/ha	vegetative growth
Horticultural in greenhouse	500-700 kg/ha	pre-sowing, pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	400-600 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	400-600 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages	300-500 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	400-600 kg/ha	pre-sowing, pre-transplant

# Cerea Blu

**11.11.16+2MgO+26SO<sub>3</sub>+0,01B+0,5Fe+0,01Zn**



Cerea Blu is an NPK mineral compound fertilizer obtained by granulation.

The nitrogen, phosphorus and potassium ratio, together with magnesium, sulphur and microelements guarantees

a balanced nutrition. The synergy between nitrogen and sulphur improves the protein content in several crops.


It is ideal for pre-sowing applications of cereal crops, oil crops and fruit trees.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>11%</b>
of which Nitrogen Ammoniacal	9,5%
of which Nitrogen Ureic	1,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>11%</b>
of which soluble in water	9%
of which soluble in neutral ammonium citrate and water	11%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>16%</b>
<b>Magnesium (MgO)</b>	<b>2%</b>
of which soluble in water	1,5%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>26%</b>
of which soluble in water	22%
<b>Boron (B)</b>	<b>0,01%</b>
<b>Iron (Fe)</b>	<b>0,5%</b>
<b>Zinc (Zn)</b>	<b>0,01%</b>

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,87 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
	300-500 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	500-700 kg/ha	post-harvest/vegetative growth
Kiwi	500-700 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	500-700 kg/ha	post-harvest/vegetative growth
Citrus trees	400-600 kg/ha	post-harvest/vegetative growth
Oive oil	500-700 kg/ha	post-harvest/vegetative growth
Table olive	500-700 kg/ha	post-harvest/vegetative growth
Extensive crops	200-300 kg/ha	localized at seeding
Horticultural in greenhouse	500-700 kg/ha	pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	300-500 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages	400-600 kg/ha	pre-transplant
Garlic, onion, leek	400-600 kg/ha	pre-sowing, pre-transplant
Alfa alfa / fodder crops	300-500 kg/ha	pre-sowing/seeding

# Super Red



Super Red is an NPK mineral compound fertilizer obtained by granulation. It contains a high amount of nitrogen in different forms (ammoniacal and ureic) and a high amount of sulphur. The synergy between nitrogen and sulphur


improves the protein content in several crops. It is indicated for pre-sowing applications on cereal crops and in post-harvest for oil crops, and fruit trees.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>14%</b>
of which Nitrogen Ammoniacal	11%
of which Nitrogen Ureic	3%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>7%</b>
of which soluble in water	6%
of which soluble in neutral ammonium citrate and water	7%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>7%</b>
<b>Calcium (CaO)</b>	<b>7%</b>
of which soluble in water	4%
<b>Magnesium (MgO)</b>	<b>2%</b>
of which soluble in water	1%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>38%</b>
of which soluble in water	35%

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,90 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Olive oil	400-600 kg/ha	post-harvest/vegetative growth
Table olive	400-600 kg/ha	post-harvest/vegetative growth
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	300-500 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	400-600 kg/ha	post-harvest/vegetative growth
Citrus trees	400-600 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	400-600 kg/ha	post-harvest/vegetative growth
Other Open field horticultural	400-600 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	300-500 kg/ha	pre-sowing, pre-transplant
Cabbages	300-500 kg/ha	pre-sowing, pre-transplant

# Terra 33



Terra 33 is an NPK mineral fertilizer obtained by granulation. It contains mineral matrices of high quality and has a slightly acidic pH that favours the mobilisation of nutrients in the soil and their absorption by the roots. The synergy between nitrogen and sulphur improves the protein content in several crops.


It is indicated for pre-sowing applications on cereal crops and fodder plants and in post-harvest for oil crops and fruit trees.



**TYPE**  
Granular



## PACKAGING



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>5%</b>
of which Nitrogen Ammoniacal	3,5%
of which Nitrogen Ureic	1,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>10%</b>
of which soluble in water	4%
of which soluble in neutral ammonium citrate and water of which soluble in formic acid	7,5%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>18%</b>
<b>Calcium (CaO)</b>	<b>11%</b>
of which soluble in water	6%
<b>Magnesium (MgO)</b>	<b>2%</b>
of which soluble in water	1,5%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>22%</b>
of which soluble in water	20%

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,95 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Legumes	200-400 kg/ha	pre-sowing/seeding
Straw cereals	200-400 kg/ha	pre-sowing/seeding
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	500-700 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	500-700 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	400-600 kg/ha	pre-sowing, pre-transplant
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruit	300-500 kg/ha	post-harvest/vegetative growth
Kiwi	400-600 kg/ha	post-harvest/vegetative growth
Wine grape	400-600 kg/ha	post-harvest/vegetative growth
Table grape	500-700 kg/ha	post-harvest/vegetative growth
Citrus trees	500-700 kg/ha	post-harvest/vegetative growth

# Bluactive



Bluactive is an NPK mineral compound fertilizer obtained by granulation with high-solubility mineral matrices. The balanced content of nitrogen, phosphorus, and potassium, together with magnesium and microelements, guarantees a complete nutrition of the crop. The high content of


potassium sulphate makes it ideal for crops that are sensitive to the presence of chlorine. Bluactive is indicated for fruit trees and a localized application at sowing (starter) on extensive crops.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>11%</b>
of which Nitrogen Ammoniacal	9,5%
of which Nitrogen Ureic	1,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>11%</b>
of which soluble in water	9%
of which soluble in neutral ammonium citrate and water	11%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>16%</b>
<b>Magnesium (MgO)</b>	<b>3%</b>
of which soluble in water	1,5%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>36%</b>
of which soluble in water	34%
<b>Boron (B)</b>	<b>0,01%</b>
<b>Zinc (Zn)</b>	<b>0,01%</b>

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,93 kg/dm<sup>3</sup>

**LOW CHLORINE**

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	300-500 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	500-700 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	500-700 kg/ha	post-harvest/vegetative growth
Nut, hazelnut , chestnut	500-700 kg/ha	post-harvest/vegetative growth
Citrus trees	400-600 kg/ha	post-harvest/vegetative growth
Oive oil	500-700 kg/ha	post-harvest/vegetative growth
Table olive	500-700 kg/ha	post-harvest/vegetative growth
Extensive crops	200-300 kg/ha	localized at seeding
Legumes	200-400 kg/ha	pre-sowing/seeding
Horticultural in greenhouse	500-700 kg/ha	pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	300-500 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages	400-600 kg/ha	pre-transplant
Garlic, onion, leek	400-600 kg/ha	pre-sowing, pre-transplant
Alfa alfa /fodder crops	300-500 kg/ha	pre-sowing/seeding

# Global



Global is an NPK mineral compound fertilizer obtained by granulation with high-solubility mineral matrices. The high content of potassium sulphate makes it ideal for crops sensitive to the presence of chlorine. The high content of soluble sulphur improves nitrogen absorp-


tion. It contains magnesium, an important element in photosynthesis processes and an essential constituent of chlorophyll. It is indicated for pre-transplant applications on horticultural crops, fruit trees and alfalfa.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>3%</b>
of which Nitrogen Ammoniacal	2%
of which Nitrogen Ureic	1%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>12%</b>
of which soluble in water	11%
of which soluble in neutral ammonium citrate and water	12%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>22%</b>
<b>Calcium (CaO)</b>	<b>6%</b>
of which soluble in water	4%
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Sulfur (SO<sub>3</sub>)</b>	<b>32%</b>
of which soluble in water	30%

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
1 kg/dm<sup>3</sup>

**LOW CHLORINE**

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Tobacco	500-700 kg/ha	pre-sowing, pre-transplant
Stone fruits	450-650 kg/ha	post-harvest/vegetative growth
Pome fruits (Apple tree, Pear tree)	350-550 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	450-650 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	450-650 kg/ha	post-harvest/vegetative growth
Citrus trees	450-650 kg/ha	post-harvest/vegetative growth
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	450-650 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	450-650 kg/ha	pre-sowing, pre-transplant
Cabbages	450-650 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	500-700 kg/ha	pre-sowing, pre-transplant

# Red Ball

## 12.6.15+6CaO+2MgO+39SO<sub>3</sub>



Red Ball is a mineral NPK complex fertilizer rich in nitrogen, potassium and sulphur to improve the quality of the productions. The elevated content of sulphur within the fertilizer acidifies the soil and allows a better assimilation


of nutrients, moreover the ratio between N/K nourishes and strengthens the plant at the same time. Magnesium stimulates photosynthesis and has a greening action. Phosphorus keeps the root system active and receptive.



**TYPE**  
Granular



**PACKAGING**



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>12%</b>
of which Nitrogen Ammoniacal	9%
of which Nitrogen Ureic	3%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>6%</b>
of which soluble in water	4%
of which soluble in neutral ammonium citrate and water	6%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>15%</b>
<b>Calcium (CaO)</b>	<b>6%</b>
of which soluble in water	3%
<b>Magnesium (MgO)</b>	<b>2%</b>
of which soluble in water	1%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>39%</b>
of which soluble in water	37%

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,92 kg/dm<sup>3</sup>

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits (pesco, nettarina, percoco, albicocco, susino, mandorlo, ciliegio)	300-500 kg/ha	post-harvest/vegetative growth
Pome fruits (Apple tree, Pear tree)	300-500 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	400-600 kg/ha	post-harvest/vegetative growth
Nut, hazelnut, chestnut	400-600 kg/ha	post-harvest/vegetative growth
Citrus trees (limone, arancio, mandarino, clementino, pompelmo)	400-600 kg/ha	vegetative growth
Small fruits (mora, ribes, lampone, mirtillo)	300-500 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	400-600 kg/ha	post-harvest/vegetative growth
Olive oil	300-500 kg/ha	vegetative growth
Table olive	300-500 kg/ha	vegetative growth
Horticultural in greenhouse (Tomato, zucchini, pear tree, melanzana, cetriolo)	500-700 kg/ha	pre-sowing, pre-transplant
Industrial crops (Tomato, Potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural (asparago, carciofo, carota, Strawberry)	400-600 kg/ha	pre-sowing, pre-transplant
Leafy vegetables (insalata, cicoria, radicchio, spinacio)	400-600 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae (Apple tree, anguria, zucca, zucchini, cetriolo)	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages (cappuccio, verza, broccolo, cavolfiore, cavoletti di Bruxelles)	300-500 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	400-600 kg/ha	pre-sowing, pre-transplant



# Base



Base is an NPK mineral compound fertilizer obtained by granulation with high-solubility matrices. The high content of phosphorus and potassium guarantees an efficient nutrition of the plant. The product is rich in soluble sulphur and the synergy between nitrogen and sulphur increases


the protein content in cereal crops. It is indicated for pre-sowing and pre-transplant applications on extensive and horticultural crops.



**TYPE**  
Granular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) total</b>	<b>6%</b>
of which Nitrogen Ammoniacal	5%
of which Nitrogen Ureic	1%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>12%</b>
of which soluble in water	11%
of which soluble in neutral ammonium citrate and water	12%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>18%</b>
<b>Calcium (CaO)</b>	<b>10%</b>
of which soluble in water	5%
<b>Sulfur (SO<sub>3</sub>)</b>	<b>34%</b>
of which soluble in water	28%

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**

0,95 kg/dm<sup>3</sup>

**LOW CHLORINE**

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Stone fruits	400-600 kg/ha	post-harvest/vegetative growth
Pome fruits	300-500 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	400-600 kg/ha	post-harvest/vegetative growth
Wine grape	300-500 kg/ha	post-harvest/vegetative growth
Table grape	400-600 kg/ha	post-harvest/vegetative growth
Citrus trees	400-600 kg/ha	post-harvest/vegetative growth
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-sowing, pre-transplant
Horticultural in greenhouse	500-700 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	500-700 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	400-600 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	400-600 kg/ha	pre-sowing, pre-transplant
Cabbages	400-600 kg/ha	pre-sowing, pre-transplant
Tobacco	500-700 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	500-700 kg/ha	pre-sowing, pre-transplant

# Mastercote (control release Nitrogen)

18.9.18+25SO<sub>3</sub>



Mastercote is an NPK mineral compound fertilizer obtained by granulation with high-solubility and availability mineral matrices. It contains 12% of coated urea which allows a controlled release of nitrogen over time and guarantees a nutrient intake in the proper phenological stages; in this

way the development of the root and the foliar systems is promoted. The high content of water-soluble sulphur helps nitrogen absorption.


It is indicated for horticultural crops, fruit trees and turf.



**TYPE**  
Granular



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>18%</b>
of which Urea ricoperta a rilascio controllato	5%
of which ureic	5,5%
of which ammoniacal	7,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) total</b>	<b>9%</b>
of which soluble in water	8,5%
of which solubile in neutral ammonium citrate and water	9%
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>18%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>25%</b>


**DENSITY**  
1,08 kg/dm<sup>3</sup>

**LOW CHLORINE**


Color and density are indicative. For hazard warnings see page 154.

## DOSAGES AND USES

CROPS	DOSAGES SPAGLIO	DOSAGES LOCALIZZATO	PERIOD
Wine grape	300-400 kg/ha	150-200 kg/ha	post-harvest/vegetative growth
Table grape	300-500 kg/ha	150-250 kg/ha	post-harvest/vegetative growth
Actinidia (kiwi)	300-500 kg/ha	150-250 kg/ha	post-harvest/vegetative growth
Citrus trees	200-400 kg/ha	100-200 kg/ha	post-harvest/vegetative growth
Stone fruits	300-400 kg/ha	150-200 kg/ha	post-harvest/vegetative growth
Pome fruits	300-400 kg/ha	150-200 kg/ha	post-harvest/vegetative growth
Nursery and garden plant	300-500 kg/ha	150-250 kg/ha	pre-sowing, pre-transplant
Turf	30-50 gr/m <sup>2</sup>	15-25 gr/m <sup>2</sup>	pre-sowing/seeding
Nut, hazelnut, chestnut	300-400 kg/ha	150 - 200 kg/ha	post-harvest/vegetative growth
Open field horticultural and in greenhouse	200-400 kg/ha	100 - 200 kg/ha	pre-sowing, pre-transplant







# ORGANIC ECO FRIENDLY AGRICULTURE

Family of fertilizers allowed in organic agriculture. Formulations from ORGANIC line are ideal for basal applications and coverage.

Organic is a family of high-quality granular fertilizers allowed in organic farming obtained by granulation. The formulations are ideal for basal and cover fertilizations. The variety of products offered allows us to guarantee the correct development of any crop while maintaining the choice of organic agriculture without loss in quality and production.

The organic nitrogen present in the Organic fertilizers ensures a slow release into the soil by microorganisms and avoids leaching, guaranteeing a prolonged availability over time throughout the entire crop cycle.

Their granulometry allows a good flexibility with applications both with fertilizer spreaders in the open field both in the hopper during sowing.

The presence of sulfur increases the protein content especially in cereal crops and improves the absorption of nitrogen in all the others limiting the losses.

# BioNascor

## 5.10+12CaO+8SO<sub>3</sub>+6Fe+12C-Org



BioNascor is an organic NP fertilizer obtained by granulation from a mixture of high quality organic matrices (bone meal, stillage, turf and hydrolyzed gelatin) allowed for organic farming. The organic matter is rich in amino acids, humic and fulvic acids which help to absorb nutrients. Phosphorus acts in the first development stages of plants; organic nitrogen is slowly released during the


whole production cycle. BioNascor is ideal for cereals, horticultural crops and fruit trees. Its granulometry allows a distribution with spreaders or micro-granulators and transplanting machines. It can be applied both in furrow and at transplant (starter technique).



**TYPE**  
MicroGranular



### PACKAGING



### COMPOSITION

Nitrogen (N) organic	5%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) total	10%
Potassium (K <sub>2</sub> O) soluble in water	1,5%
Calcium (C <sub>a</sub> O)	12%
Sulfur (SO <sub>3</sub> ) soluble in water	8%
Iron (Fe)	6%
Organic carbon	12%

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,85 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	400-600 kg/ha	pre-sowing/seeding
Extensive crops	400-600 kg/ha	pre-sowing/seeding
Legumes	400-600 kg/ha	pre-sowing/seeding
Industrial crops (Tomato, potato, sugarbeet)	300-500 kg/ha	pre-sowing, pre-transplant; localized
Alfa alfa/ fodder crops	500-700 kg/ha	pre-sowing/seeding
Open field horticultural and in greenhouse	300-500 kg/ha	pre-sowing, pre-transplant

# BioMaster

## 6.5.13+8CaO+16SO<sub>3</sub>+3Fe+20C-org

BioMaster is an NPK organo-mineral compound fertilizer allowed for organic farming and obtained by granulation of noble raw materials. The balanced content of nitrogen, phosphorus, potassium and sulphur makes it ideal for fruit trees, grapes and horticultural crops.

Its granulometry allows distributing BioMaster both with spreaders and micro-granulators (starter technique).


bioagricert 



**TYPE**  
MicroGranular



### PACKAGING



### COMPOSITION

Nitrogen (N) organic	6%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) soluble in mineral acids of which at least 55% soluble in formic acid	2%
Potassium (K <sub>2</sub> O) soluble in water	13%
Calcium (C <sub>a</sub> O)	8%
Sulfur (SO <sub>3</sub> ) soluble in water	16%
Iron (Fe)	3%
Organic carbon	20%

LOW CHLORINE

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,85 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Stone fruits	700-900 kg/ha	post-harvest/vegetative growth
Pome fruits	600-800 kg/ha	post-harvest/vegetative growth
Actinidia (Kiwi)	700-900 kg/ha	post-harvest/vegetative growth
Citrus trees	600-800 kg/ha	post-harvest/vegetative growth
Wine and table grape	600-900 kg/ha	post-harvest/vegetative growth
Olive tree	600-800 kg/ha	post-harvest/vegetative growth
Nut, hazelnut, chestnut	800-1000 kg/ha	post-harvest/vegetative growth
Straw cereals	400-600 kg/ha	pre-sowing/seeding
Extensive crops	400-600 kg/ha	pre-sowing/seeding
Legumes	400-600 kg/ha	pre-sowing/seeding
Horticultural in greenhouse	600-800 kg/ha	pre-sowing, pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	700-900 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	700-900 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	600-800 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	600-800 kg/ha	pre-sowing, pre-transplant
Cabbages	600-800 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	600-800 kg/ha	pre-sowing, pre-transplant

# BioAger

## 4.6.10+10CaO+2MgO+16SO<sub>3</sub>+3Fe+12C-Org



BioAger is an NPK organo-mineral compound fertilizer allowed for organic farming obtained by the granulation of noble raw materials. The organic matrix of BioAger has a high nutritional value and is easily absorbed by microorganisms. The organic matrices provide organic nitrogen and organic humified carbon, essential elements to improve soil structure and plant growth. It has a high content of glycine betaines, molecules that


generally develop in conditions of environmental stress; they act as osmoprotectants in plants and increase their productivity. It contains slow-release sulphur which allows a better root development. The balanced content of nitrogen, phosphorus and potassium, and the presence of calcium, magnesium and sulphur make BioAger ideal for fruit trees, Wine grape, olive trees, and horticultural crops.



**TYPE**  
MicroGranular



**PACKAGING**



**COMPOSITION**

<b>Nitrogen (N) organic</b>	<b>4%</b>
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in mineral acids of which at least 55% soluble in formic acid</b>	<b>2%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>10%</b>
<b>Calcium (C<sub>a</sub>O)</b>	<b>10%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Iron (Fe)</b>	<b>3%</b>
<b>Organic carbon</b>	<b>12%</b>

LOW CHLORINE

Color and density are indicative. For hazard warnings see page 154.

**DENSITY**  
0,85 kg/dm<sup>3</sup>

**DOSAGES AND USES**

CROPS	DOSAGES	PERIOD
Straw cereals	400-600 kg/ha	pre-sowing/seeding
Extensive crops	400-600 kg/ha	pre-sowing/seeding
Legumes	400-600 kg/ha	pre-sowing/seeding
Horticultural in greenhouse	600-800 kg/ha	pre-sowing, pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	700-900 kg/ha	pre-sowing, pre-transplant
Other Open field horticultural	700-900 kg/ha	pre-sowing, pre-transplant
Leafy vegetables	600-800 kg/ha	pre-sowing, pre-transplant
Cucurbitaceae	600-800 kg/ha	pre-sowing, pre-transplant
Cabbages	600-800 kg/ha	pre-sowing, pre-transplant
Garlic, onion, leek	600-800 kg/ha	pre-sowing, pre-transplant
Stone fruits	700-900 kg/ha	post-harvest/vegetative growth
Pome fruits	600-800 kg/ha	post-harvest/vegetative growth
Actinidia (Kiwi)	700-900 kg/ha	post-harvest/vegetative growth
Citrus trees	600-800 kg/ha	post-harvest/vegetative growth
Wine grape	600-800 kg/ha	post-harvest/vegetative growth
Table grape	700-900 kg/ha	post-harvest/vegetative growth

# Biorganic N11

11.0.0+40C-org



Biorganic is an organic nitrogen fertilizer obtained from high-quality raw materials permitted in organic farming. It contains a high level of organic nitrogen, 40% of which is water-soluble, meaning high bioavailability for soil microbial flora; furthermore, nitrogen in this form is easily mineralizable. 95% of the organic carbon contained in


Biorganic is extractable and therefore made available to soil microbial flora. The high nitrogen and carbon content makes Biorganic ideal for fertilizing cereal crops.



**TYPE**  
Flakes



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) organic</b>	<b>11%</b>
of which soluble in water	5%
<b>Organic carbon</b>	<b>40%</b>
Extractable organic carbon (C) on total organic carbon	95%

Color is indicative.

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	400-600 kg/ha	pre-seeding
Extensive crops	600-800 kg/ha	pre-seeding
Rice	400-600 kg/ha	pre semina
Other Open field horticultural	600-800 kg/ha	pre-transplant
Leafy vegetables	400-600 kg/ha	pre semina / pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	500-700 kg/ha	pre-seeding
Horticultural in green house	600-800 kg/ha	pre-transplant
Actinidia (kiwi) Kiwi	700-800 kg/ha	post-harvest
Citrus trees	700-800 kg/ha	post-harvest
Olivo tree	500-700 kg/ha	post-harvest
Wine and table grape	500-700 kg/ha	post-harvest

# Biocere 533

## 5.3.3+20C-Org




BioCere is a pelleted organo-mineral NPK fertilizer made from premium raw materials permitted in organic farming. The pellet is suitable for spreader application, and the ratio of nitrogen, phosphorus, and potassium makes the formulation particularly suitable for fertilizing fruit crops, vines, and vegetables.



**TYPE**  
Pellet



**PACKAGING**




### COMPOSITION

Nitrogen (N) organic	5%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) total	3%
Potassium (K <sub>2</sub> O) soluble in water	3%
Organic carbon	20%

Color is indicative.

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	500-600 kg/ha	pre-seeding
Extensive crops	600-700 kg/ha	pre-seeding
Rice	600-700 kg/ha	pre semina
Other Open field horticultural	800-1000 kg/ha	pre-transplant
Leafy vegetables	600-700 kg/ha	pre semina / pre-transplant
Industrial crops (Tomato, potato, sugarbeet)	600-700 kg/ha	pre-seeding
Horticultural in green house	700-800 kg/ha	pre-transplant
Actinidia (kiwi) Kiwi	600-800 kg/ha	post-harvest
Citrus trees	600-800 kg/ha	post-harvest
Olivo tree	800-1000 kg/ha	post-harvest
Wine and table grape	600-800 kg/ha	post-harvest



# SPECIALITY

## THE FUTURE IS HERE



Family of special granular fertilizers characterized by a different type of release of nutrients (more specifically for nitrogen).

Speciality line fertilizers are produced with high-quality raw materials and contain soluble and assimilable phosphorus.

The formulation of Speciality fertilizers guarantees greater agronomic efficiency with a reduction of nitrogen losses by leaching and volatilization. The different forms of nitrogen guarantee prolonged availability over time throughout the crop cycle. Furthermore, the combination of nitrogen and sulfur brings a higher protein content in cereal crops and better nitrogen absorption in all other crops limiting its losses.

The fertilizers of the Speciality family are divided into two nitrogen types:


- **Stabilized nitrogen**
- **Activated nitrogen**

Speciality nitrogen stabilized products contain a urease inhibitor (the enzyme responsible for the transformation of urea in the soil), a technology that increases urea use efficiency by limiting nitrogen volatilization losses.

Activated nitrogen products allow a gradual and progressive release of nitrogen. Activation occurs through the use of amino acids.

# THE NITROGEN CYCLE

## From fertilization to absorption by plants, much of this element does not reach its destination



Today over 46% of the nitrogen used in agriculture is distributed in the form of urea which, however, does not always manage to maximize its efficiency. When we distribute urea in the field, if in the following days there are no rains, the fertilizer remains concentrated in the first centimeters of soil and is subject to loss due to volatilization because urea is hydrolyzed in ammonia (volatile compound) and in carbon dioxide.

If this transformation takes place in the soil, in the first 15-20 cm, ammonia reacts with water and turns into ammoniacal nitrogen ( $NH_4^+$ ) retained by soil colloids. Lastly, the ammoniacal nitrogen is transformed into nitric nitrogen ( $NO_3^-$ ) by nitrifying bacteria

This form of nitrogen is the one primarily absorbed by the roots of the plants but is not retained by the soil, therefore it is subject to leaching in case of heavy rain or irrigation. In conditions of water stagnation or soil compaction, the denitrification process can be triggered which leads to the loss of nitrogen in the atmosphere ( $N_2$  and  $NO_x$ ).


Organic nitrogen is subject to mineralization processes while the mineral component can be immobilized in the organic matter depending on the soil carbon/nitrogen ratio

# Evolution 56

## 38.0.0+18SO<sub>3</sub> ACTIVATED WITH AMINO ACIDS

Evolution 56 is a granulated nitrogen-based fertilizer activated **with amino acids** obtained from enzymatic hydrolysis. It has a high sulphur content which, in synergy with nitrogen, guarantees a higher protein content in many crops. The activated formulation (urea nitrogen plus ammoniacal nitrogen) allows a gradual and progressive release of nitrogen; the ammonia fraction is already


available from the first vegetative phases, while urea nitrogen will become available in the forms assimilable by the plant at a later stage. Its formulation increases the absorption of potassium, the organoleptic qualities of plants, the efficiency of use of nutritive elements and it respects the environment.



**TYPE**  
Granular



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>38%</b>
of which Nitrogen Ureic	32%
of which Nitrogen Ammoniacal	6%
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>18%</b>

Activated with amino acids by enzymatic hydrolysis.  
Color and density are indicative. For hazard warnings see page 146.

**DENSITY**  
0,76 kg/dm<sup>3</sup>

### DOSAGES AND USES


CROPS	DOSAGES	PERIOD
Straw cereals	150-250 kg/ha	top dressing
Extensive crops	200-300 kg/ha	top dressing
Open field horticultural	200-400 kg/ha	top dressing
Citrus trees	300-400 kg/ha	da vegetative growth
Wine grape	200-300 kg/ha	vegetative growth
Table grape	300-400 kg/ha	da vegetative growth
Olive oil	200-300 kg/ha	vegetative growth
Table olive	200-300 kg/ha	vegetative growth

# Cereas 38

## 38.0.0+18SO<sub>3</sub>

Cereas 38 is a nitrogen-based fertilizer obtained by blending. It has a high content of ureic nitrogen. The combination of ureic and ammoniacal nitrogen in synergy with sulphur guarantees a high nutritional value for plants


(increase in protein content in cereal crops). Its formulation allows a maximum absorption of nitrogen. It is ideal for winter cereal crops.



**TYPE**  
Granular



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>38%</b>
of which Nitrogen Ureic	32%
of which Nitrogen Ammoniacal	6%
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>18%</b>

Color and density are indicative. For hazard warnings see page 146.

**DENSITY**  
0,76 kg/dm<sup>3</sup>

### DOSAGES AND USES


CROPS	DOSAGES	PERIOD
Straw cereals	200-300 kg/ha	top dressing
Extensive crops	250-350 kg/ha	top dressing
Open field horticultural	200-400 kg/ha	top dressing
Citrus trees	300-400 kg/ha	da vegetative growth
Wine grape	250-350 kg/ha	vegetative growth
Table grape	300-400 kg/ha	da vegetative growth
Olive oil	250-350 kg/ha	vegetative growth
Table olive	250-350 kg/ha	vegetative growth
Fruit trees	200-300 kg/ha	vegetative growth

# STABILIZED NITROGEN

Many of the nitrogen fertilizers in the SPECIALITY family feature a gradual release of nitrogen, thanks to the presence of a urease enzyme inhibitor (**NBPT**) in the soil.

This slows urea hydrolysis, resulting in reduced losses due to nitrogen volatilization.

**Slowing this process extends the time available for dissolved urea to be incorporated into the soil through rainfall or irrigation.**



The NBPT molecules replace the urea in the active site of urease (1), which will be released again only following the degradation of the NBPT itself (2-3).

The hydrolysis process, restarting (4), will allow the urea found in the soil to transform into nitrogenous forms available to the plant, reducing losses through volatilization.

## Benefits of using NBPT in our fertilizers

- 1.** Greater nitrogen efficiency (**REDUCTION OF LOSSES BY VOLATILIZATION**) with decrease in dosages and number of interventions (**MAXIMUM BENEFIT AT LOWER COST**)
- 2.** Increased production efficiency (**GREATER AVAILABILITY OF THE ELEMENT FOR THE PLANT**)
- 3.** Better vegetative-productive ratio (**STABILIZATION OF THE ELEMENT IN TIME**)
- 4.** Compared to other urease inhibitors, NBPT is **more stable** in different types of soil, **more effective** and has a **longer duration of action** under different conditions (hot and dry climate, etc.).



# CereaSlow 33


## 33.0.0+30SO<sub>3</sub> with NBPT



CereaSlow 33 is a nitrogen-based fertilizer obtained by blending with ureic nitrogen stabilised with NBPT. This is a molecule that inhibits the activity of urease enzyme in the soil, reducing losses by volatilization up to 30%. NBPT is applied to the granules through a coating process which allows to cover them uniformly, increasing their

effectiveness. The high sulphur content, in synergy with nitrogen, favours the protein synthesis processes in the plant.


It is ideal for the fertilization of cereal crops.



**TYPE**  
Granular



### PACKAGING



Color and density are indicative

### COMPOSITION

<b>Nitrogen (N) total</b>	<b>33%</b>
of which Nitrogen Ureic	23%
of which Nitrogen Ammoniacal	10%
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>30%</b>

**DENSITY**  
0,80 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	200-300 kg/ha	top dressing
Extensive crops	250-350 kg/ha	top dressing
Open field horticultural	200-400 kg/ha	top dressing
Citrus trees	300-400 kg/ha	from vegetative growth
Wine grape	250-350 kg/ha	vegetative growth
Table grape	300-400 kg/ha	from vegetative growth
Olive oil	250-350 kg/ha	vegetative growth
Table olive	250-350 kg/ha	vegetative growth
Fruit trees	200-300 kg/ha	vegetative growth

# CereaSlow 40


## 40.0.0+13SO<sub>3</sub> with NBPT



CereaSlow 40 is a nitrogen-based fertilizer obtained by blending with ureic nitrogen stabilized with NBPT. This is a molecule that inhibits the activity of urease enzyme in the soil, reducing losses by volatilization up to 30%. NBPT is applied to the granules through a coating process which allows to cover them uniformly, increasing their

effectiveness. The combination of ureic and ammoniacal nitrogen, in synergy with sulphur, favours protein synthesis processes in the plant.


It is ideal for the fertilization of cereal and extensive crops.



**TYPE**  
Granular



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>40%</b>
of which Nitrogen Ureic	35%
of which Nitrogen Ammoniacal	5%
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>13%</b>

Color and density are indicative

**DENSITY**  
0,75 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	200-300 kg/ha	top dressing
Extensive crops	250-350 kg/ha	top dressing
Open field horticultural	200-400 kg/ha	top dressing
Citrus trees	250-350 kg/ha	from vegetative growth
Wine grape	250-350 kg/ha	vegetative growth
Table grape	250-350 kg/ha	from vegetative growth
Olive oil	250-350 kg/ha	vegetative growth
Table olive	250-350 kg/ha	vegetative growth
Fruit trees	200-300 kg/ha	vegetative growth

# CereaSlow 46


## 46.0.0 with NBPT



CereaSlow 46 is a nitrogen-based fertilizer obtained by blending ureic nitrogen stabilized with NBPT. This is a molecule that inhibits the activity of urease enzyme in the soil, reducing losses by volatilization up to 30%. NBPT

is applied to the granules through a coating process which allows to cover them uniformly, increasing their effectiveness.


It is ideal for the fertilization of cereal crops.



**TYPE**  
Granular



### PACKAGING



### COMPOSITION

**Nitrogen (N) Ureic** **46%**

Color and density are indicative

**DENSITY**  
0,70 kg/dm<sup>3</sup>

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	150-250 kg/ha	top dressing
Extensive crops	250-400 kg/ha	top dressing
Open field horticultural	200-350 kg/ha	top dressing
Citrus trees	250-350 kg/ha	from vegetative growth
Wine grape	200-300 kg/ha	vegetative growth
Table grape	200-300 kg/ha	from vegetative growth
Olive oil	200-300 kg/ha	vegetative growth
Table olive	200-300 kg/ha	vegetative growth
Fruit trees	200-300 kg/ha	vegetative growth




# StarSlow

## 30.15+2MgO with NBPT

StarSlow is an NP granular fertilizer obtained by blending ureic nitrogen stabilized with NBPT. This is a molecule that inhibits the activity of urease enzyme in the soil, reducing losses by volatilization up to 30%. NBPT is applied to the granules through a coating process which allows to cover


them uniformly, increasing their effectiveness. StarSlow has a high phosphorus content which is water-soluble and easy to be absorbed by the plant. It is ideal for pre-sowing or localized applications.



**TYPE**  
Granular



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>30%</b>
of which Nitrogen Ureic	24%
of which Nitrogen Ammoniacal	6%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in neutral ammonium citrate and water</b>	<b>15%</b>
of which soluble in water	14%
<b>Magnesium (MgO)</b>	<b>2%</b>

Color and density are indicative

**DENSITY**  
0,94 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	250-300 kg/ha	pre-sowing
Extensive crops	300-350 kg/ha	pre-sowin / mechanical weed
Open field horticultural	250-350 kg/ha	pre-sowing, pre-transplant


# Più Sprint

## 24.9.5+5CaO+2MgO+10SO<sub>3</sub> with NBPT



Più Sprint is a nitrogen fertilizer obtained by blending ureic nitrogen stabilized with NBPT. This molecule inhibits the urease activity of the soil, preventing volatilization losses and reducing them by up to 30%. NBPT is applied to the granule through a coating process that allows the granules


to be uniformly covered, increasing their use efficiency. The high water solubility of the other components combined with the high quality of potassium makes this product extremely effective in a wide range of applications..



**TYPE**  
Granular



### PACKAGING



### COMPOSITION

<b>Nitrogen (N) total</b>	<b>24%</b>
of which Nitrogen Ureic	20,5%
of which Nitrogen Ammoniacal	3,5%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in neutral ammonium citrate and water</b>	<b>9%</b>
of which soluble in water	8%
<b>Potassium (K) soluble in water</b>	<b>5%</b>
<b>Sulfur (SO<sub>3</sub>) soluble in water</b>	<b>10%</b>
<b>Calcium (CaO) soluble in water</b>	<b>5%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Iron (Fe) soluble in water</b>	<b>0,02%</b>
<b>Manganese (Mn) soluble in water</b>	<b>0,01%</b>
<b>Zinc (Zn) soluble in water</b>	<b>0,002%</b>

Color and density are indicative

**DENSITY**  
0,98 kg/dm<sup>3</sup>

### DOSAGES AND USES

CROPS	DOSAGES	PERIOD
Straw cereals	200-250 kg/ha	top dressing
Extensive crops	200-300 kg/ha	pre-sowing / mechanical weed
Open field horticultural	200-400 kg/ha	top dressing
Stone fruits	350-450 kg/ha	vegetative growth
Pome fruits	300-400 kg/ha	vegetative growth
Nut, hazelnut , chestnut	300-400 kg/ha	vegetative growth
Actinidia (kiwi)	350-450 kg/ha	vegetative growth
Citrus trees	300-400 kg/ha	vegetative growth
Wine grape	300-400 kg/ha	vegetative growth
Table grape	350-450 kg/ha	vegetative growth
Olive oil	300-400 kg/ha	vegetative growth
Table olive	350-450 kg/ha	vegetative growth




# WATER SOLUBLE

A hand is shown pouring water from a clear glass into a small potted plant. The background is a blurred greenhouse or nursery setting with other plants. The entire image has a teal overlay.

# FERTIGATION

Water-soluble fertilizer



# FERTIGATION

## WATER-SOLUBLE FERTILIZER



FERTIGATION line is characterized by low salinity products suitable for the various phenological phases of cultivation and that can be applied both in fertigation and leaf treatments.

The absence of chlorine, sodium and other substances harmful to the plant makes them particularly suitable for different types of use.

Fertigation fertilizers have the ability to lower the soil acidity and they are highly soluble.

# Fertigation 20.20.20

Fertigation 20.20.20 is a water-soluble NPK crystalline fertilizer that can be applied in fertigation.

The balanced nitrogen, phosphorus, and potassium ratio makes it available for all stages of the crop cycle by providing a balanced nutrition. Nitrogen is present in three forms (nitric, ammoniacal, ureic) that ensure a constant release throughout the whole crop cycle. The high phosphorus content improves the root system.

It contains microelements chelated with EDTA, that guarantee an easier assimilation preventing possible deficiencies and favouring biochemical processes.


It is particularly indicated for horticultural crops and fruit trees.



**TYPE**  
Cristallino



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>20%</b>
of which ammoniacal	4%
of which ureic	10,1%
of which nitrico	5,9%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>20%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>20%</b>
<b>Boron (B) soluble in water</b>	<b>0,02%</b>
<b>Iron (Fe) chelated EDTA</b>	<b>0,07%</b>
<b>Manganese (Mn) chelated EDTA</b>	<b>0,03%</b>
<b>Molibdeno (Mo) soluble in water</b>	<b>0,003%</b>
<b>Zinc (Zn) chelated EDTA</b>	<b>0,05%</b>

Color and density are indicative. For hazard warnings see page 146.

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIOD
Stone fruits	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Pome fruits	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Citrus trees	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Wine grape	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Table grape	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Olive tree	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Nut, hazelnut , chestnut	30-50 kg/ha	250-300 g/hl	All fasi vegetative
Horticultural in greenhouse	3,5-5 kg/1000 m2	200-250 g/hl	All fasi vegetative
Open field horticultural	30-50 kg/ha	250-300 g/hl	All fasi vegetative

# Fertigation 10.40.10

Fertigation 10.40.10 is a water-soluble NPK crystalline fertilizer that can be applied in fertigation. The high phosphorus content helps the root development of the plant and the first growth stages after transplant. Magnesium helps to improve physiological processes, such


as photosynthesis. It contains microelements chelated with EDTA, that ensure an easier assimilation preventing possible deficiencies and favouring biochemical processes.



**TYPE**  
Cristallino



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>10%</b>
of which ammoniacal	7,8%
of which nitrico	2,2%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>40%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>10%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Iron (Fe) chelated EDTA</b>	<b>0,03%</b>
<b>Manganese (Mn) chelated EDTA</b>	<b>0,01%</b>
<b>Molibdeno (Mo) soluble in water</b>	<b>0,001%</b>
<b>Zinc (Zn) chelated EDTA</b>	<b>0,02%</b>

Color and density are indicative

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIODO
Stone fruits	30-50 kg/ha	250-300 g/hl	vegetative growth, swelling and hardening of the hazel
Pome fruits	30-50 kg/ha	250-300 g/hl	from vegetative growth to flowering
Citrus trees	30-50 kg/ha	250-300 g/hl	from vegetative growth to flowering
Wine grape	30-50 kg/ha	250-300 g/hl	from vegetative growth to flowering
Table grape	30-50 kg/ha	250-300 g/hl	from vegetative growth to flowering
Olive tree	30-50 kg/ha	250-300 g/hl	from vegetative growth to flowering
Nut, hazelnut , chestnut	30-50 kg/ha	250-300 g/hl	from vegetative growth to flowering
Horticultural in greenhouse	3,5-5 kg/1000 m2	200-250 g/hl	from post-transplant
Open field horticultural	30-50 kg/ha	250-300 g/hl	from post-transplant

# Fertigation 15.5.26

Fertigation 15.5.26 is a water-soluble NPK crystalline fertilizer. The high potassium content supports the plant at fruit set. Magnesium helps to improve some physiological processes in the plant, such as photosynthesis. It contains microelements chelated with EDTA, that ensure an easier

assimilation preventing possible deficiencies and favouring biochemical processes.


It can be applied in fertigation.



**TYPE**  
Cristallino



## PACKAGING



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>15%</b>
of which ammoniacal	5,4%
of which ureic	2,6%
of which nitrico	7%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>5%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>26%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Iron (Fe) chelated EDTA</b>	<b>0,03%</b>
<b>Manganese (Mn) chelated EDTA</b>	<b>0,01%</b>
<b>Molibdeno (Mo) soluble in water</b>	<b>0,001%</b>
<b>Zinc (Zn) chelated EDTA</b>	<b>0,02%</b>

**LOW CHLORINE**

Color and density are indicative

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIODO
Stone fruits	30-50 kg/ha	250-300 g/hl	from veraison
Pome fruits	30-50 kg/ha	250-300 g/hl	from veraison
Citrus trees	30-50 kg/ha	250-300 g/hl	from fruit swelling
Wine grape	30-50 kg/ha	250-300 g/hl	from veraison
Table grape	30-50 kg/ha	250-300 g/hl	from veraison
Olive tree	30-50 kg/ha	250-300 g/hl	from veraison
Nut, hazelnut , chestnut	30-50 kg/ha	250-300 g/hl	from fruit swelling
Horticultural in greenhouse	3,5-5 kg/1000 m2	200-250 g/hl	from veraison
Open field horticultural	30-50 kg/ha	250-300 g/hl	from veraison
Leafy vegetables	30-50 kg/ha	250-300 g/hl	from half cycle

# Fertigation 17.6.21

Fertigation 17.6.21 is a water-soluble NPK crystalline fertilizer. The high nitrogen and potassium formulation supports the plant during fruit set. Thanks to the different forms of nitrogen (nitric, ammoniacal, ureic), it guarantees a constant intake of the nutrient throughout the whole crop cycle.


Calcium helps to prevent some physiopathologies such as blossom-end rot in tomato/pepper/melon, tip burn in lettuce, bitter-pit in apple trees. It contains microelements chelated with EDTA, that ensure an easier assimilation preventing possible deficiencies and favouring biochemical processes. It can be applied in fertigation.



**TYPE**  
Cristallino



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>17%</b>
of which ammoniacal	0,6%
of which ureic	5,8%
of which nitrico	10,6%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>6%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>21%</b>
<b>Calcium (C<sub>2</sub>O) soluble in water</b>	<b>8%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Iron (Fe) chelated EDTA</b>	<b>0,03%</b>
<b>Manganese (Mn) chelated EDTA</b>	<b>0,01%</b>
<b>Molibdeno (Mo) soluble in water</b>	<b>0,001%</b>
<b>Zinc (Zn) chelated EDTA</b>	<b>0,02%</b>

Color and density are indicative. For hazard warnings see page 154.

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIODO
Stone fruits	30-50 kg/ha	250-300 g/hl	from veraison
Pome fruits	30-50 kg/ha	250-300 g/hl	from veraison
Citrus trees	30-50 kg/ha	250-300 g/hl	from fruit swelling
Wine grape	30-50 kg/ha	250-300 g/hl	from veraison
Table grape	30-50 kg/ha	250-300 g/hl	from veraison
Olive tree	30-50 kg/ha	250-300 g/hl	from veraison
Nut, hazelnut , chestnut	30-50 kg/ha	250-300 g/hl	from fruit swelling
Horticultural in greenhouse	3,5-5 kg/1000 m <sup>2</sup>	200-250 g/hl	from veraison
Open field horticultural	30-50 kg/ha	250-300 g/hl	from veraison
Leafy vegetables	30-50 kg/ha	250-300 g/hl	from half cycle

# Fertigation 7.15.30

Fertigation 7.15.30 is a water-soluble NPK crystalline fertilizer. The high phosphorus and potassium formulation makes it ideal for the fertilization of horticultural crops before transplant. Magnesium helps to improve physiological processes of the plant, such as photosynthesis. It contains microelements chelated


with EDTA, that ensure an easier assimilation preventing possible deficiencies and favouring biochemical processes. It can be applied in fertigation.



**TYPE**  
Cristallino



**PACKAGING**



## COMPOSITION

<b>Nitrogen (N) total</b>	<b>7%</b>
of which ammoniacal	1,8%
of which ureic	0,8%
of which nitrico	4,4%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>15%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>30%</b>
<b>Magnesium (MgO) soluble in water</b>	<b>2%</b>
<b>Boron (B) soluble in water</b>	<b>0,01%</b>
<b>Iron (Fe) chelated EDTA</b>	<b>0,03%</b>
<b>Manganese (Mn) chelated EDTA</b>	<b>0,01%</b>
<b>Molibdeno (Mo) soluble in water</b>	<b>0,001%</b>
<b>Zinc (Zn) chelated EDTA</b>	<b>0,02%</b>

Color and density are indicative

LOW CHLORINE

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIODO
Stone fruits	30-50 kg/ha	250-300 g/hl	from unripe fruit
Pome fruits	30-50 kg/ha	250-300 g/hl	from unripe fruit
Citrus trees	30-50 kg/ha	250-300 g/hl	from unripe fruit
Wine grape	30-50 kg/ha	250-300 g/hl	from bunch elongation
Table grape	30-50 kg/ha	250-300 g/hl	from bunch elongation
Olive tree	30-50 kg/ha	250-300 g/hl	from unripe fruit
Nut, hazelnut , chestnut	30-50 kg/ha	250-300 g/hl	from unripe fruit
Horticultural in greenhouse	3,5-5 kg/1000 m <sup>2</sup>	200-250 g/hl	from unripe fruit
Open field horticultural	30-50 kg/ha	200-250 g/hl	from unripe fruit
Leafy vegetables	30-50 kg/ha	200-250 g/hl	from half cycle

# Fertigation 30.10.10

Fertigation 30.10.10 is a water-soluble NPK crystalline fertilizer. The high nitrogen content makes it ideal during the vegetative development stages before blossoming. Thanks to the different forms of nitrogen (nitric, ammoniacal, ureic), it guarantees a constant intake of the


nutrient throughout the whole crop cycle. It contains microelements chelated with EDTA, that ensure an easier assimilation preventing possible deficiencies and favouring the biochemical processes of the plant. It can be applied in fertigation.



**TYPE**  
Cristallino



**PACKAGING**



## COMPOSITION


<b>Nitrogen (N) total</b>	<b>30%</b>
of which ammoniacal	4,2%
of which ureic	22,8%
of which nitrico	3%
<b>Phosphorus (P<sub>2</sub>O<sub>5</sub>) soluble in water</b>	<b>10%</b>
<b>Potassium (K<sub>2</sub>O) soluble in water</b>	<b>10%</b>
<b>Boron (B) soluble in water</b>	<b>0,02%</b>
<b>Iron (Fe) chelated EDTA</b>	<b>0,07%</b>
<b>Manganese (Mn) chelated EDTA</b>	<b>0,03%</b>
<b>Molibdeno (Mo) soluble in water</b>	<b>0,003%</b>
<b>Zinc (Zn) chelated EDTA</b>	<b>0,05%</b>

Color and density are indicative. For hazard warnings see page 154.

**LOW CHLORINE**

## DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIODO
Stone fruits	30-50 kg/ha	250-300 g/hl	vegetative development
Pome fruits	30-50 kg/ha	250-300 g/hl	vegetative development
Citrus trees	30-50 kg/ha	250-300 g/hl	vegetative development
Wine grape	30-50 kg/ha	250-300 g/hl	from 5cm sprout to visible cluster
Table grape	30-50 kg/ha	250-300 g/hl	from 5cm sprout to visible cluster
Olive tree	30-50 kg/ha	250-300 g/hl	all vegetative stages
Nut, hazelnut, chestnut	30-50 kg/ha	250-300 g/hl	vegetative development
Horticultural in greenhouse	3,5-5 kg/1000 m <sup>2</sup>	200-250 g/hl	first vegetative stages
Open field horticultural	30-50 kg/ha	200-250 g/hl	first vegetative stages
Leafy vegetables	30-50 kg/ha	200-250 g/hl	post-transplant to half cycle



# HAZARD WARNINGS

## SPECIAL NUTRITION

<b>B-Power</b>	<b>H360FD</b> Suspected of damaging fertility or the unborn child. Restricted to professional users.	
<b>Calciomagno</b>	<b>H302:</b> Harmful if swallowed; <b>H318:</b> Causes serious eye damage; <b>EUH071:</b> Corrosive to the respiratory tract	
<b>Calcito</b>	<b>H314:</b> Causes severe skin burns and eye damage	
<b>Color MgZn</b>	<b>H411:</b> Toxic to aquatic life with long lasting effects; <b>H319:</b> Causes serious eye irritation	
<b>Crisco</b>	<b>H317:</b> May cause an allergic skin reaction; <b>H412:</b> Harmful to aquatic life with long lasting effects	
<b>Edafos</b>	<b>H302:</b> Harmful if swallowed; <b>H314:</b> Causes severe skin burns and eye damage	
<b>Febo Bio</b>	<b>H360FD:</b> Suspected of damaging fertility or the unborn child; <b>H373:</b> May cause damage to organs through prolonged or repeated exposure; <b>H318:</b> Causes serious eye damage; <b>H411:</b> Toxic to aquatic life with long lasting effects	
<b>Febo Mix</b>	<b>H360FD:</b> Suspected of damaging fertility or the unborn child; <b>H317:</b> May cause an allergic skin reaction; <b>H318:</b> Causes serious eye damage; <b>H411:</b> Toxic to aquatic life with long lasting effects	
<b>Focus Ca</b>	<b>H302:</b> Harmful if swallowed; <b>H314:</b> Causes severe skin burns and eye damage	
<b>Glycos Plus</b>	<b>H360FD:</b> Suspected of damaging fertility or the unborn child; Restricted to professional users	
<b>Hydrostar</b>	<b>EUH210:</b> Safety data sheet available on request	
<b>Leaf K</b>	<b>EUH210:</b> Safety data sheet available on request	
<b>Leaf P-Ca</b>	<b>H302:</b> Harmful if swallowed; <b>H314:</b> Causes severe skin burns and eye damage	
<b>MagnetiCal</b>	<b>H319:</b> Toxic to aquatic life with long lasting effects	
<b>Leaf S-Quality</b>	<b>EUH210:</b> Safety data sheet available on request	
<b>Nano.T Fe</b>	<b>H314:</b> Causes severe skin burns and eye damage	
<b>Nano.T Fe Bio</b>	<b>H314:</b> Causes severe skin burns and eye damage	
<b>Nano.T Cu 4.5</b>	<b>H302:</b> Harmful if swallowed; <b>H315:</b> Causes skin irritation; <b>H319:</b> Causes serious eye irritation; <b>H400:</b> Very toxic to aquatic life; <b>H411:</b> Toxic to aquatic life with long lasting effects	

<b>Nano.T Cu Bio</b>	<b>H318:</b> Causes serious eye damage; <b>H410:</b> Molto Toxic to aquatic life with long lasting effects	 
<b>Nano.T Cu-Zn</b>	<b>H318:</b> Causes serious eye damage; <b>H319:</b> Causes serious eye irritation; <b>H400:</b> Very toxic to aquatic life; <b>H411:</b> Toxic to aquatic life with long lasting effects	 
<b>Nano.T. CaPO</b>	<b>H319:</b> Causes serious eye irritation	
<b>Nano.T Carbo</b>	<b>H315:</b> Causes skin irritation; <b>H319:</b> Causes serious eye irritation	
<b>Nano.T Total</b>	<b>H314:</b> Causes severe skin burns and eye damage	
<b>Nano.T Zn</b>	<b>H318:</b> Causes serious eye damage; <b>H411:</b> Toxic to aquatic life with long lasting effects	 
<b>Oro-B</b>	<b>H360FD:</b> Suspected of damaging fertility or the unborn child;	
<b>Proser Ca</b>	<b>H314:</b> Causes severe skin burns and eye damage	
<b>Proser MnZn</b>	<b>H318:</b> Causes serious eye damage; <b>H410:</b> Molto Toxic to aquatic life with long lasting effects	 
<b>Verv Plus</b>	<b>H319:</b> Causes serious eye irritation; <b>H412:</b> Harmful to aquatic life with long lasting effects <b>H360FD</b> Suspected of damaging fertility or the unborn child. Restricted to professional users.	 
<b>VigorGreen</b>	<b>H360FD</b> Suspected of damaging fertility or the unborn child. Restricted to professional users.	

## GRANULAR

<b>ACTIVE ACTIVE PREMIUM</b>	<b>H318:</b> Causes serious eye damage	
<b>FERT FERT PREMIUM</b>	<b>H318:</b> Causes serious eye damage	
<b>NUECR4</b>	<b>H318:</b> Causes serious eye damage	
<b>ORGANIC</b>	<b>H315:</b> Causes skin irritation; <b>H319:</b> Causes serious eye irritation	

## WATER-SOLUBLE

<b>Fertigation NPK 17.6.21</b>	<b>H302:</b> Harmful if swallowed; <b>H318:</b> Causes serious eye damage	 
<b>Fertigation NPK 20.20.20</b>	<b>EUH210:</b> Safety data sheet available on request	
<b>Fertigation NPK 30.10.10</b>	<b>EUH210:</b> Safety data sheet available on request	







**PRODUCTS CATALOG**

**CERE A FCP**

Via Farfusola 6, 37050  
Bonavicina di S. Pietro di Morubio (VR) - Italy  
Tel. +39.045.7125911  
fpcerea@fpcerea.it - www.fpcerea.it

1-2026-3

