

Fertigation 30.10.10

Fertigation 30.10.10 is a water-soluble NPK crystalline fertilizer. The high nitrogen content makes it ideal during the vegetative development stages before blossoming. Thanks to the different forms of nitrogen (nitric, ammoniacal, ureic), it guarantees a constant intake of the


nutrient throughout the whole crop cycle. It contains microelements chelated with EDTA, that ensure an easier assimilation preventing possible deficiencies and favouring the biochemical processes of the plant. It can be applied in fertigation.



TYPE
Cristallino



PACKAGING



COMPOSITION

Nitrogen (N) total	30%
of which ammoniacal	4,2%
of which ureic	22,8%
of which nitrico	3%
Phosphorus (P₂O₅) soluble in water	10%
Potassium (K₂O) soluble in water	10%
Boron (B) soluble in water	0,02%
Iron (Fe) chelated EDTA	0,07%
Manganese (Mn) chelated EDTA	0,03%
Molibdeno (Mo) soluble in water	0,003%
Zinc (Zn) chelated EDTA	0,05%

Color and density are indicative. For hazard warnings see page 154.

LOW CHLORINE

DOSAGES AND USES

CROPS	FERTIGATION	FOLIAR G/HL	PERIODO
Stone fruits	30-50 kg/ha	250-300 g/hl	vegetative development
Pome fruits	30-50 kg/ha	250-300 g/hl	vegetative development
Citrus trees	30-50 kg/ha	250-300 g/hl	vegetative development
Wine grape	30-50 kg/ha	250-300 g/hl	from 5cm sprout to visible cluster
Table grape	30-50 kg/ha	250-300 g/hl	from 5cm sprout to visible cluster
Olive tree	30-50 kg/ha	250-300 g/hl	all vegetative stages
Nut, hazelnut , chestnut	30-50 kg/ha	250-300 g/hl	vegetative development
Horticultural in greenhouse	3,5-5 kg/1000 m2	200-250 g/hl	first vegetative stages
Open field horticultural	30-50 kg/ha	200-250 g/hl	first vegetative stages
Leafy vegetables	30-50 kg/ha	200-250 g/hl	post-transplant to half cycle