

Giove Bio Gold

100% vegetable amino acids

bioagricert 



GIOVE BIO GOLD has a very high concentration of vegetal amino acids (98%) in medium-short chains with a phytostimulant, anti-stress and nutritional action. GIOVE BIO GOLD is effective at low doses thanks to the highly soluble powder formulation. GIOVE BIO GOLD is rapidly absorbed and activates the plant's metabolism allowing for optimal development and rapid recovery of stressed and/or blocked plants.

Benefits


- High effectiveness at low dosages thanks to the very high concentration;
- Biostimulating action thanks to the high presence of peptides, oligopeptides and free amino acids, all of vegetal origin;
- Activation of plant metabolism thanks to the high content of totally soluble organic nitrogen;
- Environmentally friendly and innovative as it comes from the enzymatic hydrolysis of plant tissues;
- Easy to use thanks to the high solubility;
- Increased effectiveness of the products distributed together thanks to the co-formulating action of the amino acid matrix.



TYPE

Water-soluble powder

PACKAGING



COMPOSITION

Nitrogen (N) organic	16%
Carbon (C) organic	50%
pH	6,5

ALSO CONTAINS:

Total amino acids	98%
of which Glicina	25,5%
of which Prolina	14%

Color and density are indicative

DENSITY

0,60 +/- 0,05 kg/dm³

DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	1-2 kg/ha	vegetative growth, before blossoming, after fruit set, fruit development	foliar
Wine and table grape	1-2k g/ha	pre-blossoming, blossoming, berries groat-sized, bunches begin to hang, berries swelling	foliar
Olive tree	1-2 kg/ha	pre-blossoming, blossoming, cluster flowers, drupe swelling	foliar
HazelNut	2 kg/ha	from fruit set	foliar
Open field horticultural (Potato, Tomato, carciofo...)	1-3 kg/ha	post-transplant/ post emergence, vegetative development, post fruit set, fruit development	foliar
Horticultural in greenhouse	300-500 g/1000m ²	3-4applications 10-15 days after transplant	foliar
Cereals	1-2 kg/ha	tillering, leaf flag	foliar
Potato	1 kg/ha	10-15 days after emergence until tuberification	foliar
All crops	5-6 kg/ha	2-4 applications	fertigation