

# Mycovix

## Resilient Roots



MYCOVIX has a high content of specific microorganisms that, thanks to their intense activity, are able to colonize the soil rapidly. This characteristic creates an ideal environment for root development (production of hormone-like substances, production of organic acids, and improvement of the soil's biological value) and an unfavorable environment for phytopathogens (space and Nutrient competition).

MYCOVIX is an innovative fertilizer that promotes the development of a healthy rhizosphere and stimulates the plant's natural defenses, improving its resistance, resilience, and vitality. MYCOVIX works synergistically to enhance the

growth and robustness of crops, helping maintain a healthy and vibrant soil ecosystem.

### Benefits

- Promotes the growth of beneficial microorganisms such as *Pochonia c.* and *Purpureocillium l.*;
- Increased root development;
- More fertile soil;
- Improved plant health;
- Enhanced Nutrient absorption.



**TYPE**  
Liquid



### PACKAGING



TRANSPARENT



### COMPOSITION

Mycorrhizae	0,1%
Rhizosphere bacteria (CFU/g)	2x10 <sup>7</sup>
Trichoderma	1x10 <sup>4</sup>
pH	7

### DENSITY

1,13 +/- 0,05 kg/dm<sup>3</sup>

The product does not contain genetically modified organisms and pathogenic organisms. Color and density are indicative

## DOSAGES AND USES

CROPS	DOSAGES	PERIOD	MODE
Fruit trees	3-4 l/ha	1-2 spring applications 1-2 summer applications	fertigation
Open field horticultural	3-5 l/ha	From post-transplant (2-3 applications every 8-10 gg)	fertigation
Horticultural in greenhouse	400-500ml/1000m	From post-transplant every 8-12 gg	fertigation

## DIRECTION FOR USE

PREBIOTIC	DO NOT MIX WITH	NOTES
SuprEmo, Giove Bio Gold, Giove Bio, Verv N9, Naturblackk	Stillages, phosphoric acid, nitric acid, products containing sulfuric acid, products containing copper, sulfur, bactericidal products, phosphites, and fungicides.	Space treatments with products containing phosphorus at least 7 days apart. Space any treatments with phosphites, fungicides, and copper-based products at least 10-15 days apart.