



Tomato

Managing two key phases in tomato cultivation: transplant and ripening

Be-Start NP applied during transplant or in the following phases allows the plant to **overcome stress from transplant more rapidly** and to promptly react to non optimal weather conditions (late frosts, low or excessive rainfalls).

Calcito and **Proser MnZn** combined with calcium nitrate stimulate root activity **improving nutrient absorption**, in particular calcium. The plant will get more compact, reducing waste (green and rotten tomatoes) and **increasing commercial harvest**.

The early application of **Proser Ca**, from the stage of fruit set of the first truss, **reduces the risk of apical rot**.

Nano.T Fe prevents iron chlorosis.

Nano.t Cu increases the resistance of the crop to biotic and abiotic stresses.



**Grow well
to eat better**

Technical line

GRANULAR

Cereaphos 250-300 kg/ha
or **Universal Up 250-300 kg/ha**
Soil fertilization



FERTIGATION

Be-Start 5.15 15 l/ha

Transplant

Leaf P-Ca 10 l/ha + Nano.T Fe 5 l/ha

After 7 days

Leaf P-Ca 10 l/ha -

After 10-15 days from transplant

Calcito 20 l/ha* + 25 kg/ha Nitrato di Calcio + Proser MnZn 2 l/ha

Second application after 20 days – fruit set of the first truss

Calcito 10 l/ha* + 50 kg/ha Nitrato di Calcio + Proser MnZn 1 l/ha

Third application after 20 days – fruit set of the third truss



To improve the absorption of fertilizers it is recommended to employ at each application **Verv N9 15 l/ha**

** in saline, sodium, calcareous ("white" soils) increase the doses of Calcite by 10 l/ha*

FOLIAR APPLICATIONS

Giove Bio Gold 1 kg/ha + Leaf P-Ca 2 l/ha

Transplant

Nano.T Cu 1,5 l/ha + Leaf P-Ca 2 l/ha


After 7 days

Proser Ca 2 l/ha


Fruit set of the first truss



Results of three field trials carried out in Verona in 2020-21



Results obtained applying 15 kg/ha of Be-Start NP at transplant and after 7 days



Results obtained with three applications of Calcito, Calcium Nitrate and Proser MnZn as indicated above in the application period and dosage

for more details: agronomia@fpcerea.it

www.fpcerea.it

