


Fertigation

20.20.20




FORMULATION



Title and composition

Nitrogen (N)	Phosphorus (P ₂ O ₅)	Potassium (K ₂ O)	Copper EDTA (Cu)	Boron (B)	Iron EDTA (Fe)	Manganese EDTA (Mn)	Molybdenum (Mo)	Zinc EDTA (Zn)
20%	20%	20%	0.003%	0.02%	0.07%	0.03%	0.003%	0.05%

PACKAGING



TIPOLOGY

CRYSTALLINE



Dosages and uses

Crop	Dosages	Period and method of use
citrus fruits	20-50 kg/ha	every 10-15 days from fruit set to veraison
stone fruits	40-100 kg/ha	every 10-15 days from fruit set to veraison
kiwi	20-50 kg/ha	every 10-15 days from fruit set to veraison
olive tree	20-50 kg/ha	every 10-15 days from fruit set to veraison
open field horticultural	40-100 kg/ha	every 10-15 days from post-transplant
greenhouse horticultural	4-8 kg / 1.000 m ²	every 10-15 days from post-transplant
pome fruits	40-100 kg/ha	every 10-15 days from fruit set to veraison
grapevine	40-100 kg/ha	every 10-15 days from fruit set to veraison

To be used in fertigation, apply according to the needs of the crop.

The Fertigation products in the low chlorine content version can be used not only in fertigation, but also by foliar application.

The products are perfectly soluble in water.

For foliar application we recommend 2-4 kg/ha diluted in 400 L of water.