



NANO.T

# NANO.T® Cu

## High Efficiency Copper


NANO.T Cu 4.5 contains copper with high nutritional efficiency thanks to the small size (nanotechnology) of the particles present in the formulation which makes it effective even at low dosage. NANO.T Cu 4.5 is ideal for TREATING DEFICIENCIES AND PREVENTING PHYSIOLOGICAL and mechanical DAMAGES (cracks, hail, pruning and harvesting) of the foliar and root system of the plant. NANO.T Cu 4.5, being a colloidal suspension of nano particles, has a wide contact surface and it does not leach when applied on the leaf and on the soil. NANO.T Cu 4.5 strengthens the plant and plant tissues and stimulates self-defense mechanisms to improve the response to disease and nutritional disorders.



### Benefits

- Low-dosage effectiveness thanks to the wide contact surface;
- Low environmental impact as it does not leach if applied on the leaf and in fertigation;
- Low risk of phytotoxicity
- It doesn't stain crops thanks to the innovative formulation;
- It improves the effectiveness of the combined products (whether they are fertilizers and / or pesticides), by acidifying the solution.

### FORMULATION



## Title and composition

Copper (Cu)	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	Potassium (K <sub>2</sub> O)	Sulphur (SO <sub>3</sub> )	pH
4,5% (1) 2,5% (3) 2% (2)	3,0%	4,0% (2)	7,0% (2)	3,3

(1) Total - (2) Water soluble - (3) nano form

### PACKAGING



## Dosages and uses

Crop	Dosages	Application
Pome fruit (apple, pear)	1-2 l/ha	foliar
Stone fruit (Peach, Nectarine, Percoco, Apricot, Plum, Almond, Cherry)	1-1,5 l/ha	foliar
Olive (for oil and table). Kiwi, Walnut	3-4 l/ha	foliar
Wine grapes and table grapes	2-4 l/ha	foliar
Citrus fruits (Orange, Mandarin, Clementine, Lemon, Grapefruit, Cedar)	3-3,5 l/ha	foliar
Hazelnut, Chestnut	3-3,5 l/ha	foliar
Vegetables (processing tomato, table tomato, pepper, aubergine, courgette, cucumber, pumpkin, melon, watermelon, strawberry, artichoke)	3-3,5 l/ha	foliar
Potato, Carrot, Onion, Garlic, Leek, Beetroot	2-3 l/ha	foliar
Spinach, leafy vegetables (lettuce, radicchio, chicory)	2-2,5 l/ha	foliar
Broccoli, Cabbage, Cauliflower, Fennel	1-1,5 l/ha	foliar
Small fruits (blueberry, raspberry, blackberry, etc.)	1,5-2 l/ha	foliar
All crops	2,5-3 l/ha	fertigation

### TIPOLOGY

LIQUID

