


# Fertigation

30.10.10


Fertigation 30.10.10 is a water-soluble NPK crystalline fertilizer. The high nitrogen content makes it ideal during the vegetative development stages before blossoming. Thanks to the different forms of nitrogen (nitric, ammoniacal, ureic), it guarantees a constant intake of the nutrient throughout the whole crop cycle. It contains microelements chelated with EDTA, that ensure an easier assimilation preventing possible deficiencies and favouring the biochemical processes of the plant. It can be applied in fertigation.



## Title and composition

| Nitrogen (N) | Phosphorus (P <sub>2</sub> O <sub>5</sub> ) | Potassium (K <sub>2</sub> O) | Copper EDTA (CU) | Boron (B) | Iron EDTA (Fe) | Manganese EDTA (Mn) | Molybdenum (Mo) | Zinc EDTA (Zn) |
|--------------|---|------------------------------|------------------|-----------|----------------|---------------------|-----------------|----------------|
| 30%          | 10%   | 10%                          | 0,003%           | 0,02%     | 0,07%          | 0,03%               | 0,003%          | 0,05%          |
| Low chlorine |   |                              |                  |           |                |                     |                 |                |

## FORMULATION



## Dosages and uses

| Crop  | Fertigation Dosages          | Foliar Dosages | Period and method of use           |
|---|------------------------------|----------------|------------------------------------|
| Stone fruit (peach, nectarine, cherry, etc.)  | 30-50 kg/ha                  | 250-300 g/hl   | vegetative development             |
| Pome fruit (apple, pear)                      | 30-50 kg/ha                  | 250-300 g/hl   | vegetative development             |
| Citrus fruits (lemon, orange, mandarin, etc.) | 30-50 kg/ha                  | 250-300 g/hl   | vegetative development             |
| Grapevine                                     | 30-50 kg/ha                  | 250-300 g/hl   | from 5cm sprout to visible cluster |
| Table grape                                   | 30-50 kg/ha                  | 250-300 g/hl   | from 5cm sprout to visible cluster |
| Olive tree                                    | 30-50 kg/ha                  | 250-300 g/hl   | all vegetative stages              |
| Walnut, hazelnut, chestnut                    | 30-50 kg/ha                  | 250-300 g/hl   | vegetative development             |
| Horticultural in greenhouse                   | 3,5-5 kg/1000 m <sup>2</sup> | 200-250 g/hl   | first vegetative stages            |
| Open field horticultural                      | 30-50 kg/ha                  | 200-250 g/hl   | first vegetative stages            |
| Leafy vegetables                              | 30-50 kg/ha                  | 200-250 g/hl   | post-transplant to half cycle      |

## PACKAGING



## TYPOLOGY

CRYSTALLINE



For foliar application we recommend 2-4 kg/ha diluted in 400 L of water.